

User Manual for the LMD Martian Atmospheric General Circulation Model

Ehouarn MILLOUR, François FORGET
*Contributors to earlier versions: Karine Dassas,
Christophe HOURDIN, Frédéric HOURDIN,
and Yann WANHERDRICK*
initial version translated by Gwen Davis
(LMD)

April 4, 2017

Contents

1	Introduction	4
2	Main features of the model	5
2.1	Basic principles	5
2.2	Dynamical-Physical separation	5
2.3	Grid boxes:	6
2.3.1	Horizontal grids	6
2.3.2	Vertical grids	8
2.4	Variables used in the model	10
2.4.1	Dynamical variables	10
2.4.2	Physical variables	10
2.4.3	Tracers	10
3	The physical parameterizations of the Martian model: some references	12
3.1	General	12
3.2	Radiative transfer	12
3.2.1	CO₂ gas absorption/emission:	12
3.2.2	Absorption/emission and diffusion by dust:	13
3.3	Subgrid atmospheric dynamical processes	13
3.3.1	Turbulent diffusion in the upper layer	13
3.3.2	Convection	14
3.3.3	Effects of subgrid orography and gravity waves	14
3.4	Surface thermal conduction	14
3.5	CO ₂ Condensation	14
3.6	Tracer transport and sources	14
3.7	Thermosphere	15
4	Running the model: a practice simulation	16
4.1	Obtaining the model	16
4.2	Prerequisites	17
4.3	Installing the model	17
4.3.1	Settings in LMDZ.MARS (deprecated)	17
4.3.2	Settings in LMDZ.COMMON (advised)	18
4.4	Compiling the model	18
4.4.1	Compiling in LMDZ.MARS (deprecated)	18
4.4.2	Compiling in LMDZ.COMMON (advised)	19
4.5	Input files (initial states and def files)	19
4.6	Running the model	20
4.7	Visualizing the output files	20
4.7.1	Using GrAds to visualize outputs	20
4.8	Resuming a simulation	22
4.9	Chain simulations	22

4.10	Creating and modifying initial states	23
4.10.1	Using program “newstart”	23
4.10.2	Creating the initial start_archive.nc file	24
4.10.3	Changing the horizontal or vertical grid resolution	24
5	Program organization and compilation script	25
5.1	Organization of the model source files	26
5.2	Programming	26
5.3	Model organization	26
5.4	Compiling the model	26
6	Input/Output	29
6.1	NetCDF format	29
6.1.1	NetCDF file editor: ncdump	29
6.1.2	Graphic visualization of the NetCDF files using GrAds	30
6.2	Input and parameter files	30
6.2.1	run.def	31
6.2.2	callphys.def	33
6.2.3	traceur.def	36
6.2.4	z2sig.def	36
6.2.5	Initialization files: start and startfi	37
6.3	Output files	43
6.3.1	NetCDF restart files - restart.nc and restartfi.nc	43
6.3.2	NetCDF file - diagfi.nc	43
6.3.3	Stats files	44
7	Zoomed simulations	48
7.1	To define the zoomed area	48
7.2	Making a zoomed initial state	48
7.3	Running a zoomed simulation and stability issue	49
8	Water Cycle Simulation	50
9	Photochemical Module	52
10	1D version of the Mars model	54
10.1	Compilation	54
10.2	1-D runs and input files	54
10.3	Output data	56
A	GCM Martian Calendar	57
B	Utilities	59
B.1	concatnc	59
B.2	lslin	59
B.3	localtime	59
B.4	zrecast	59
B.5	lslin	59
B.6	hrecast	60
B.7	expandstartfi	60
B.8	extract	60

Chapter 1

Introduction

This document is a user manual for the General Circulation Model of the Martian atmosphere developed by the Laboratoire de Météorologie Dynamique of the CNRS in Paris in collaboration with the Atmospheric and Oceanic Planetary Physics sub-department in Oxford. It corresponds to the version of the model available since November 2002, that includes the new dynamic code lmdz3.3 and the input and output data in NetCDF format. The physical part has been available since June 2001, including the NLTE radiative transfer code valid at up to 120 km, tracer transport, the water cycle with water vapour and ice, the "double mode" dust transport model, and with optional photochemistry and extension in the thermosphere up to 250km.

A more general, scientific description of the model without tracers can be found in *Forget et al.* [1999].

Chapter 2 of this document, to be read before any of the others, describes the main features of the model. The model is divided into two relatively independent parts: (1) The hydrodynamic code, that is shared by all atmospheres (Earth, Mars, etc.) that integrates the fluid mechanics equations in time and on the globe, and (2) a set of Martian physical parameterizations, including, for example, the radiative transfer calculation in the atmosphere and the turbulence mix in the upper layer.

It is followed by a list of references for anyone requiring a detailed description of the physics and the numerical formulation of the parameterizations of the Martian physical part (Chapter 3).

For your **first contact with the model**, chapter 4 guides the user through a practice simulation (choosing the initial states and parameters and visualizing the output files).

The document then describes the programming code for the model, including a user computer manual for compiling and running the model (Chapter 5).

Chapter 6 describes the input/output data of the model. The input files are the files needed to initialize the model (state of the atmosphere at the initial time t_0 as well as a dataset of boundary conditions) and the output files are "time series", i.e. records of the atmospheric flow evolution as simulated by the model, the "diagfi files", the "stats files", the daily averages etc. Some means to edit or visualize these files (editor "ncdump" and the graphics software "grads") are also described.

Chapter 8 explains how to run a simulation including the water cycle. Chapter 9 illustrates how to run the model with the photochemical module.

Finally, chapter 10 will help you to use a 1-dimensional version of the model, which may be a simpler tool for some analysis work.

Chapter 2

Main features of the model

2.1 Basic principles

The General Circulation Model (GCM) calculates the temporal evolution of the different **variables** (listed below) that control or describe the Martian meteorology and climate at different points of a **3D “grid”** (see below) that covers the entire atmosphere.

From an initial state, the model calculates the evolution of these variables, timestep by timestep:

- At instant t , we know variable X_t (temperature for example) at one point in the atmosphere.
- We calculate the evolution (the **tendencies**) $(\frac{\partial X}{\partial t})_1$, $(\frac{\partial X}{\partial t})_2$, etc. arising from each physical phenomenon, calculated by a **parameterization** of each of these phenomenon (for example, heating due to absorption of solar radiation).
- At the next time step $t + \delta t$, we can calculate $X_{t+\delta t}$ from X_t and $(\frac{\partial X}{\partial t})$. This is the **“integration”** of the variables in time. (For example, $X_{t+\delta t} = X_t + \delta t(\frac{\partial X}{\partial t})_1 + \delta t(\frac{\partial X}{\partial t})_2 + \dots$)

The main task of the model is to calculate these tendencies $(\frac{\partial X}{\partial t})$ arising from the different parameterized phenomenon.

2.2 Dynamical-Physical separation

In practice, the 3D model operates in two parts:

- one **dynamical part** containing the numerical solution of the general equations for atmospheric circulation. This part (including the programming) is common to the Earth and Martian model, and in general for all atmospheres of the terrestrial type.
- a second **physical part** that is specific to the planet in question and which calculates the forced circulation and the climate details at each point.

The calculations for the dynamical part are made on a 3D grid with horizontal exchanges between the grid boxes, whereas the physical part can be seen as a juxtaposition of atmosphere “columns” that do not interact with each other (see diagram 2.1).

The dynamical and physical parts deal with variables of different natures, and operate on grids that are differently constructed. The temporal integration of the variables is based on different numerical schemes (simple, such as the one above for the physical part, and more complicated, the “Matsuno-Leapfrog” scheme for the dynamical part). The timesteps are also different. The physical timestep is `iphysiq` times longer than the dynamical




Figure 2.1: Physical/dynamical interface

timestep, as the solution of the dynamic equations requires a shorter timestep than the forced calculation for the physical part.

In practice, the main program that handles the whole model (`gcm.F`) is located in the dynamical part. When the temporal evolution is being calculated, at each timestep the program calls the following:

1. Call to the subroutine that handles the total tendency calculation ($\frac{\partial X}{\partial t}$) arising from the dynamical part (`caldyn.F`)
2. Integration of these dynamical tendencies to calculate the evolution of the variables at the following timesteps (subroutine `integrd.F`)
3. Every `iphysiq` dynamical timestep, a call to the interface subroutine (`calfis.F`) with the physical model (`physiq.F`), that calculates the evolution of some of the purely physical variables (e.g: surface temperature `tsurf`) and returns the tendencies ($\frac{\partial X}{\partial t}$) arising from the physical part.
4. Integration of the physical variables (subroutine `addfi.F`)
5. Similarly, calculation and integration of tendencies due to the horizontal dissipation and the “sponge layer” is done every `idissip` dynamical time step.

Remark: The physical part can be run separately for a 1-D calculation for a single column using program `testphys1d.F`.

2.3 Grid boxes:

Examples of typical grid values are `64x48x25`, `64x48x32` or `32x24x25` in longitudelatitudealtitude. For Mars (radius~3400 km), a `64x48` horizontal grid corresponds to grid boxes of the order of `330x220` kilometers near the equator.

2.3.1 Horizontal grids

Dynamics and physics use different grids. Figure 2.2 shows the correspondance and indexing of the physical and dynamical grids as well as the different locations of variables on these grids. To identify the coordinates of a variable (at one grid point up, down, right or left) we use coordinates `r lonu`, `r latu`, `r lonv`, `r latv` (longitudes and latitudes, in radians).

On the dynamical grid, values at `i=1` are the same as at `i=IM+1` as the latter node is a redundant point (due to the periodicity in longitude, these two nodes are actually located




Figure 2.2: Dynamical and physical grids for a 6×7 horizontal resolution. In the dynamics (but not in the physics) winds u and v are on specific staggered grids. Other dynamical variables are on the dynamical "scalar" grid. The physics uses the same "scalar" grid for all the variables, except that nodes are indexed in a single vector containing $NGRID=2+(JM-1) \times IM$ points when counting from the north pole.
 N.B.: In the Fortran program, the following variables are used: $iim=IM$, $iip1=IM+1$, $jjm=JM$, $jjp1=JM+1$.

at the same place). Similarly, the extreme $j=1$ and $j=JM+1$ nodes on the dynamical grid (respectively corresponding to North and South poles) are duplicated $IM+1$ times. In contrast, the physical grid does not contain redundant points (only one value for each pole and no extra point along longitudes), as shown in figure 2.2. In practice, computations relative to the physics are made for a series of `ngrid` atmospheric columns, where $NGRID=IM \times (JM-1)+2$.

2.3.2 Vertical grids




Figure 2.3: Sketch illustrating the difference between hybrid and non-hybrid coordinates

The GCM was initially programmed using sigma coordinates $\sigma = p/p_s$ (atmospheric pressure over surface pressure ratio) which had the advantage of using a constant domain ($\sigma = 1$ at the surface and $\sigma = 0$ at the top of the atmosphere) whatever the underlying topography. However, it is obvious that these coordinates significantly disturb the stratospheric dynamical representation as the topography is propagated to the top of the model by the coordinate system. This problem can elegantly be solved by using a hybrid sigma-P (sigma-pressure) hybrid coordinate which is equivalent to using σ coordinates near the surface and gradually shifting to purely pressure p coordinates with increasing altitude. Figure 2.3 illustrates the importance of using these hybrid coordinates compared to simple σ coordinates. The distribution of the vertical layers is irregular, to enable greater precision at ground level. In general we use 25 levels to describe the atmosphere to a height of 80 km, 32 levels for simulations up to 120 km, or 50 levels to rise up to thermosphere. The first layer describes the first few meters above the ground, whereas the upper layers span several kilometers. Figure 2.4 describes the vertical grid representation and associated variables.

DYNAMICS		PHYSICS
-----		-----
[coordinates ap(),bp()]		[pressures]
ap(11m+1)=0,bp(11m+1)=0	*****	plev(nlayer+1)=0
aps(11m),bps(11m)	.. 11m nlayer ...	play(nlayer)
ap(11m),bp(11m)	*****	plev(nlayer)
aps(11m-1),bps(11m-1)	.. 11m-1 nlayer-1 .	play(nlayer-1)
ap(11m-1),bp(11m-1)	*****	plev(nlayer-1)
	:	:
	:	:
	:	:
aps(2),bps(2)	... 2 2	play(2)
ap(2),bp(2)	*****	plev(2)
aps(1),bps(1)	... 1 1	play(1)
ap(1)=1,bp(1)=0	*****surface*****	plev(1)=Ps

Figure 2.4: Vertical grid description of the 11m (or nlayer) atmospheric layers in the programming code (11m is the variable used in the dynamical part, and nlayer is used in the physical part). Variables ap, bp and aps, bps indicate the hybrid levels at the interlayer levels and at middle of the layers respectively. Pressure at the interlayer is $Plev(l) = ap(l) + bp(l) \times Ps$ and pressure in the middle of the layer is defined by $Play(l) = aps(l) + bps(l) \times Ps$, (where Ps is surface pressure). Sigma coordinates are merely a specific case of hybrid coordinates such that $aps = 0$ and $bps = P/Ps$. Note that for the hybrid coordinates, $bps = 0$ above ~ 50 km, leading to purely pressure levels. The user can choose whether to run the model using hybrid coordinates or not by setting variable hybrid in run.def to True or False.

2.4 Variables used in the model

2.4.1 Dynamical variables

The dynamical state variables are the atmospheric temperature, surface pressure, winds and tracer concentrations. In practice, the formulation selected to solve the equations in the dynamics is optimised using the following less “natural” variables:

- **potential temperature** θ (teta in the code), linked to temperature \mathbf{T} by $\theta = T(P/Pref)^{-\kappa}$ with $\kappa = R/C_p$ (note that κ is called kappa in the dynamical code, and r_{cp} in the physical code). We take $Pref = 610$ Pa on Mars.
- **surface pressure** (p_s in the code).
- **mass** the atmosphere mass in each grid box (*masse* in the code).
- **the covariant meridional and zonal winds** u_{cov} and v_{cov} . These variables are linked to the “natural” winds by $u_{cov} = c_u * u$ and $v_{cov} = c_v * v$, where c_u and c_v are constants that only depend on the latitude.
- **mixing ratio of tracers** in the atmosphere, typically expressed in kg/kg (array q in the code).

u_{cov} and v_{cov} , “vectorial” variables, are stored on “scalari” grids u and v respectively, in the dynamics (see section 2.2). $teta$, q , p_s , *masse*, “scalar variables”, are stored on the “scalar” grid of the dynamics.

2.4.2 Physical variables

In the physics, the state variables of the dynamics are transmitted via an interface that interpolates the winds on the scalar grid (that corresponds to the physical grid) and transforms the dynamical variables into more “natural” variables. Thus we have winds \mathbf{u} and \mathbf{v} ($m.s^{-1}$), temperature \mathbf{T} (K), pressure at the middle of the layers **play** (Pa) and at interlayers **plev** (Pa), tracers \mathbf{q} , etc. (kg/kg) on the same grid.

Furthermore, the physics also handle the evolution of the purely physical state variables:

- **co2ice** CO₂ ice on the surface ($kg.m^{-2}$)
- **tsurf** surface temperature (K),
- **tsoil** temperature at different layers under the surface (K),
- **emis** surface emissivity,
- **q2** wind variance, or more precisely the square root of the turbulent kinetic energy.
- **qsurf** tracer on the surface ($kg.m^{-2}$).

2.4.3 Tracers

The model may include different types of tracers:

- dust particles, which may have several modes
- chemical species which depict the chemical composition of the atmosphere
- water vapor and water ice particles
- ...

In the code, all tracers are stored in one three-dimensional array q , the third index of which corresponds to each individual tracer. In input and output files (“start.nc”, “startfi.nc”, see Section 4) tracers are stored separately using their individual names. Loading specific tracers requires that the appropriate tracer names are set in the `traceur.def` file (see Section 6.2.3), and specific computations for given tracers (e.g.: computing the water cycle, chemistry in the upper atmosphere, ...) is controlled by setting corresponding options in the `callphys.def` file (see Section 6.2.2).

Chapter 3

The physical parameterizations of the Martian model: some references

3.1 General

The Martian General Circulation Model uses a large number of physical parameterizations based on various scientific theories and some generated using specific numerical methods.

A list of these parameterizations is given below, along with the most appropriate references for each one. Most of these documents can be consulted at: <http://www-mars.lmd.jussieu.fr/mars/publi.html>.

General references:

A document attempts to give a complete scientific description of the current version of the GCM (a version without tracers):

- *Forget et al.* [1999] (article published in the JGR)

3.2 Radiative transfer

The radiative transfer parameterizations are used to calculate the heating and cooling ratios in the atmosphere and the radiative flux at the surface.

3.2.1 CO₂ gas absorption/emission:

Thermal IR radiation

(`lwmain`)

- New numerical method, solution for the radiative transfer equation: *Dufresne et al.* [2005].
- Model validation and inclusion of the “Doppler” effect (but using an old numerical formulation): *Hourdin* [1992] (article).
- At high altitudes, parameterization of the thermal radiative transfer (`nltecool`) when the local thermodynamic balance is no longer valid (e.g. within 0.1 Pa) : Lopez-Valverde et al. [2001] : Report for the ESA available on the web as: “CO₂ non-LTE cooling rate at 15-um and its parameterization for the Mars atmosphere”.

Absorption of near-infrared radiation

(nirco2abs)

- *Forget et al.* [1999]

3.2.2 Absorption/emission and diffusion by dust:

Dust spatial distribution

(aeropacity)

- The method for semi-interactive dust vertical distribution is detailed in *Madeleine et al.* [2011]
- Vertical distribution and description of “MGS” and “Viking” scenarios in the ESA report *Mars Climate Database V3.0 Detailed Design Document* by Lewis et al. (2001), available on the web.
- For the “MY24”-“MY26” scenarios, the dust distributions were derived from observations made by TES data is used. See technical note WP12.2.1 of ESA contract Ref ESA 11369/95/NL/JG(SC) ”New dust scenarios for the Mars Climate Model : Martian Years 24-29”, available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp12.1.1.pdf>

Thermal IR radiation

(lwmain)

- Numerical method: *Toon et al.* [1989]
- Optical properties of dust: *Madeleine et al.* [2011]

Solar radiation

(swmain)

- Numerical method: *Toon et al.* [1989]
- Optical properties of dust: see the discussion in *Madeleine et al.* [2011], which quotes properties from *Wolff et al.* [2009].

3.3 Subgrid atmospheric dynamical processes

3.3.1 Turbulent diffusion in the upper layer

(vdifc)

- Implicit numerical scheme in the vertical: see the thesis of Laurent Li (LMD, Université Paris 7, 1990), Appendix C2.
- Calculation of the turbulent diffusion coefficients: *Forget et al.* [1999].
- fluxes in the near-surface layer: *Colaitis et al.* [2012], technical note WP13.1.3d of ESA contract Ref ESA 11369/95/NL/JG(SC) ”New Mars Climate Model: d) New convection and boundary layer schemes and their impact on Mars meteorology”, available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp13.1.3d.pdf>

3.3.2 Convection

(convadj)

- For some details on the convective adjustment, see *Hourdin et al.* [1993]
- The thermals' mass flux scheme is described in *Colaitis et al.* [2012], technical note WP13.1.3d of ESA contract Ref ESA 11369/95/NL/JG(SC) "New Mars Climate Model: d) New convection and boundary layer schemes and their impact on Mars meteorology", available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp13.1.3d.pdf>

3.3.3 Effects of subgrid orography and gravity waves

(calldrag_noro , drag_noro)

See *Forget et al.* [1999] and *Lott and Miller* [1997]

3.4 Surface thermal conduction

(soil)

The numerical scheme is described in section 2 of technical note WP11.1 of ESA contract Ref ESA 11369/95/NL/JG(SC) "Improvement of the high latitude processes in the Mars Global Climate Model", available online at http://www-mars.lmd.jussieu.fr/WP2008/Polar_processes.pdf

3.5 CO₂ Condensation

- In *Forget et al.* [1998] (article published in *Icarus*):
 - Numerical method for calculating the condensation and sublimation levels at the surface and in the atmosphere (newcondens) explained in the appendix.
 - Description of the numerical scheme for calculating the evolution of CO₂ snow emissivity (co2snow) explained in section 4.1
- Noncondensable gaz treatment: see *Forget et al.* [2008], available online at <http://www.lpi.usra.edu/meetings/modeling2008/pdf/9106.pdf>
- Inclusion of sub-surface water ice table thermal effect, varying albedo of polar caps and tuning of the CO₂ cycle are described in technical note WP13.1.3e of ESA contract Ref ESA 11369/95/NL/JG(SC) "New Mars Global Climate Model: e) Improved CO₂ cycle and seasonal pressure variations", available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp13.1.3e.pdf>

3.6 Tracer transport and sources

- "Van-Leer" transport scheme used in the dynamical part (tracv1 and vlsplt in the dynamical part): *Hourdin and Armengaud* [1999]
- Transport by turbulent diffusion (in vdifc), convection (in convadj), sedimentation (sedim), dust lifting by winds (dustlift): see note "Preliminary design of dust lifting and transport in the Model" (ESA contract, Work Package 4, 1998, available on the web).
- Dust transport by the "Mass mixing ratio / Number mixing ratio" method for grain size evolution: see article by *Madeleine et al.* [2011]

- **Watercycle**, see *Montmessin et al.* [2004] and technical note WP13.1.3c of ESA contract Ref ESA 11369/95/NL/JG(SC) "New Mars Climate Model: c) Inclusion of cloud microphysics, dust scavenging and improvement of the water cycle", available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp13.1.3c.pdf>
- Radiative effect of clouds: see technical note WP13.1.3b of ESA contract Ref ESA 11369/95/NL/JG(SC) "New Mars Climate Model: b) Radiative effects of water ice clouds and impact on temperatures", available online at <http://www-mars.lmd.jussieu.fr/WP2011/wp13.1.3b.pdf>
- **Chemistry**, see *Lefèvre et al.* [2004] and *Lefèvre et al.* [2008]

3.7 Thermosphere

- A general description of the model is given in *González-Galindo et al.* [2009]
- Details on photochemistry and EUV radiative transfer can be found in *Angelats i Coll et al.* [2005] and *González-Galindo et al.* [2005]

Chapter 4

Running the model: a practice simulation

This chapter is meant for first time users of the LMD model. As the best introduction to the model is surely to run a simulation, here we explain how to go about it. All you will need are files and scripts necessary to build the GCM (all are in the `LMDZ.COMMON` and `LMDZ.MARS` directories which you will download as explained in the next sections) as well as some initial states to initiate simulations and, if not working on the LMD machines, some extra datafiles for external forcings (topography, dust scenarios, radiative properties of dust and water ice, etc.).

Once you have followed the example given below, you can then go on to change the control parameters and the initial states as you wish. A more detailed description of the model's organization as well as associated inputs and outputs are given in sections 5 and 6.

4.1 Obtaining the model

The LMD model project is developed using subversion (svn), the free software versioning and a revision control system. To obtain (download) the latest version of the model, simply go to the directory where you want to install the model and use the relevant svn command:
`svn checkout http://svn.lmd.jussieu.fr/Planeto/trunk --depth empty`

Then move to the newly generated `trunk` directory and download (i.e. update) the `LMDZ.MARS` and `LMDZ.COMMON` directories (the contents of these directories are described in chapter 5).

If you are not on the LMD machines, you will also need to download a set of files available online at:

`http://www.lmd.jussieu.fr/~lmdz/planets/mars/datadir`
(preserve the file names and subdirectory structure). This directory contains input files (topography, dust scenarios, radiative properties of scatterers, etc.) which the GCM needs to run. Where you put your local `datadir` directory (or whatever name you choose for this directory) is not critical, as that location can be specified at runtime (see sections 4.6 and 6.2.2).

To run the model, you will also need some initial condition files that can be downloaded from:

`http://www.lmd.jussieu.fr/~lmdz/planets/mars/starts` (see section 4.5).

4.2 Prerequisites

Before downloading and installing the model, a few prerequisites must be satisfied:

1. The NetCDF library must be installed on your system, using the same compiler suite (e.g. gfortran and gcc, or ifort and icc) that you will use to compile the model. The latest version of the NetCDF package is available on the web at the following address:

```
http://www.unidata.ucar.edu/software/netcdf
```

along with instructions for building (or downloading precompiled binaries of) the library.

2. Some software to load and display NetCDF files such as GrAdS or Ferret should be installed on your system.
3. The fcm utility must be installed on your system. If it is not already so, it may be obtained by the following svn command:

```
svn checkout http://forge.ipsl.jussieu.fr/fcm/svn/PATCHED/FCM.V1.2
```

And add its bin subdirectory to your PATH environment variable to make the fcm command available from anywhere.

4.3 Installing the model

There are two distinct ways of installing/compiling the model: One (nowdays deprecated) using scripts in the LMDZ.MARS directory, and the other (encouraged because one can then also run the model on parallel computers) in the LMDZ.COMMON directory

4.3.1 Settings in LMDZ.MARS (deprecated)

- Set some environment variables needed for the compilation of the model (it is also possible to set the environment variables in the makegcm script, as explained below):

LMDGCM : Path to the directory where you have put the model (full path).

If using Csh:

```
setenv LMDGCM /where/you/put/the/model/LMDZ.MARS
```

If using Bash:

```
export LMDGCM=/where/you/put/the/model/LMDZ.MARS
```

LIBOGCM : Path to the directory (libo for example) where intermediate objects will be stored during the compilation of the model with the makegcm script (if that directory does not exist then makegcm will create it).

If using Csh:

```
setenv LIBOGCM /where/you/want/objects/to/go/libo
```

If using Bash:

```
export LIBOGCM=/where/you/want/objects/to/go/libo
```

- Knowing where your NetCDF library is installed, set environment variables **NCDFINC** and **NCDFLIB**:

Once the NetCDF library has been compiled (or downloaded), you should have access to the library `libnetcdf.a` itself, the various files (`netcdf.inc`, `netcdf.mod`, ...) to include in programs, and basic NetCDF software (`nc-dump` and `ncgen`).

To ensure that during compilation, the model can find the NetCDF library and include files, you must declare environment variables **NCDFLIB** and **NCDFINC** (again, it is also possible to set these environment variables in the `makegcm` script, as explained below).

NCDFLIB must contain the path to the directory containing the object library `libnetcdf.a` and **NCDFINC** must contain the path to the directory containing the include files (`netcdf.inc`,...)

If using Csh:

```
setenv NCDFINC /wherever/is/netcdf/include
setenv NCDFLIB /wherever/is/netcdf/lib
```

If using Bash:

```
export NCDFINC=/wherever/is/netcdf/include
export NCDFLIB=/wherever/is/netcdf/lib
```

- Go to your `LMDZ.MARS` and adapt the `makegcm` script to fit your needs:
 - Examples of `makegcm` scripts, adapted for different compilers (`pgf90`, `g95`, `gfortran` and `ifort`) are provided (files `makegcm`, `makegcm_g95`, `makegcm_gfortran`, `makegcm_ifort`) copy or rename the relevant one as `makegcm` in the same directory.
 - As mentioned above, you may edit the script to hard code values of `LMDGCM`, `LIBOGCM`, `NCDFINC` and `NCDFLIB` instead of relying on the use of environment variables (see the commented out examples in the scripts at lines 20-30). Note that since the `makegcm` is a Csh script, Csh syntax must be used there.

4.3.2 Settings in `LMDZ.COMMON` (advised)

4.4 Compiling the model

4.4.1 Compiling in `LMDZ.MARS` (deprecated)

As explained above, compiling the model is done using the `makegcm` script and providing it with the appropriate options:

- Example 1: Compiling the Martian model at grid resolution `64x48x25` for example, type (in compliance with the manual for the `makegcm` function given in section 5.4)

```
makegcm -d 64x48x25 -p mars gcm
```

You will find executable **gcm.e** (the compiled model) in the directory where you ran the `makegcm` command.

- Example 2: Compiling the the Martian model with your choice of compiler options, e.g. to check for array overflow (useful for debugging: warning, the model is then much slower!):

```
makegcm -d 64x48x25 -p mars -O "-C" gcm
```

Note that the `makegcm` script also has a "debug" option which includes a collection of adequate debugging options. To use it, simply add the `-debug` option:

```
makegcm -d 64x48x25 -p mars -debug gcm
```

4.4.2 Compiling in LMDZ.COMMON (advised)

The Bash script `makelmdz_fcm` is used to compile the model. It needs not be modified or adapted to your settings, as all specificities are set in corresponding files located in the `arch` subdirectory. For a given machine, e.g. `MyMachine`, one should create two files, `arch-MyMachine.fcm` and `arch-MyMachine.path`, using the provided example files to set appropriate compiler options and paths (for instance `arch-linux-ifort-para.fcm` and `arch-linux-ifort-para.path` are adapted to run on local LMD machines).

The `makelmdz_fcm` script has the mandatory option `-arch MyArch` to specify the arch files to use (the "MyArch" string should be replaced with the name used for your own arch files), and multiple options:

- Example 1: Compiling the Martian model at grid resolution 64x48x32

```
makelmdz_fcm -arch linux-ifort -d 64x48x29 -p mars gcm
```

The executable, `gcm_64x48x29_phymars_seq.e` in the present case, will be generated in the `bin` subdirectory.

- Example 2: Compiling as above but in "debug" mode

```
makelmdz_fcm -arch linux-ifort -d 64x48x29 -p mars -debug gcm
```

- Example 3: Compiling the model to run in parallel (MPI) mode:

```
makelmdz_fcm -arch linux-ifort -parallel mpi -d 64x48x29 -p mars gcm
```

- For an overview of all available options:

```
makelmdz_fcm -h
```

4.5 Input files (initial states and def files)

In directory `LMDZ.MARS/defTank` you will find some examples of run parameter files (`.def` files) which the model needs at runtime. The four files the model requires (they must be in the same directory as the executable `gcm.e`) are: **run.def** (described in section 6.2) **callphys.def** (see section 6.2.2), **callphys.def**, **z2sig.def** and **traceur.def**.

The example `.def` files given in the `defTank` directory are for various configurations (e.g. model resolution), copy (and eventually rename these files to match the generic names) to the directory where you will run the model.

Copy initial condition files **start.nc** and **startfi.nc** (described in section 6.2) to the same directory.

You can extract such files from **start_archive** 'banks of initial states' (i.e. files which contain collections of initial states from standard scenarios and which can thus be used to check if the model is installed correctly) stored on the LMD website at <http://www.lmd.jussieu.fr/~lmdz/planets/mars/starts>. See section 4.10 for a description of how to proceed to extract **start** files from **start_archives**.

4.6 Running the model

Once you have the program `gcm.e`, input files `start.nc` `startfi.nc`, and parameter files `run.def` `callphys.def` `traceur.def` `z2sig.def` in the same directory, simply execute the program to run¹ a simulation:

```
gcm.e
```

You might also want to keep all messages and diagnostics written to standard output (i.e. the screen). You should then redirect the standard output (and error) to some file, e.g. `gcm.out`:

```
gcm.e > gcm.out 2>&1
```

4.7 Visualizing the output files

As the model runs it generates output files `diagfi.nc` and `stats.nc` files. The former contains instantaneous values of various fields and the later statistics (over the whole run) of some variables.

4.7.1 Using GrAds to visualize outputs

If you have never used the graphic software **GrAds**, we strongly recommend spending half an hour to familiarize yourself with it by following the demonstration provided for that purpose. The demo is fast and easy to follow and you will learn the basic commands. To do this read file

```
/distrib/local/grads/sample
```

For example, to visualize files `diagfi.nc` and `stats.nc`

NetCDF files `diagfi.nc` and `stats.nc` can be accessed directly using GrAdS thanks to utility program `gradsnc`, (the user does not need to intervene).

To visualize the temperature in the 5th layer using file `diagfi.nc` for example:

- GrAdS session:

```
grads return
return (opens a landscape window)
ga-> sdfopen diagfi.nc
ga-> query file (displays info about the open file, including the name of the
stored variables. Shortcut: q file)
ga-> set z 5 (fixes the altitude to the 5th layer)
ga-> set t 1 (fixes the time to the first stored value)
ga-> query dims (indicates the fixed values for the 4 dimensions. Shortcut: q
dims)
ga-> display temp (displays the temperature card for the 5th layer and for
the first time value stored. Shortcut: d T)
ga-> clear (clears the display. Shortcut: c)
```

¹Note that if you are not running on the LMD machines, you'll have to modify or add, in file `callphys.def`, the line: `datadir = /path/to/datafile`
Where `/path/to/datafile` is the full path to the directory which contains the set of files downloaded from: <http://www.lmd.jussieu.fr/~lmdz/planets/mars/datadir+>




Figure 4.1: Input/output data

```

ga-> set gxout shaded (not a contour plot, but a shaded one)
ga-> display temp
ga-> set gxout contour (returns to contour mode to display the levels)
ga-> display temp (superimposes the contours if the clear command is not
used)

```

4.8 Resuming a simulation

At the end of a simulation, the model generates **restart** files (files `restart.nc` and `restartfi.nc`) which contain the final state of the model. As shown in figure 4.1, these files (which are of the same format as the start files) can later be used as initial states for a new simulation.

The **restart** files just need to be renamed:

```

mv restart.nc start.nc
mv restartfi.nc startfi.nc

```

and running a simulation with these will in fact resume the simulation from where the previous run ended.

4.9 Chain simulations

In practice, we recommend running a chain of simulations lasting several days or longer (or hundreds of days at low resolution).

To do this, a script named `run0` is available in `LMDZ.MARS/deftank`, which should be used as follows:

- Set the length of each simulation in `run.def` (i.e. set the value of `nday`)
- Set the maximum number of simulations at the beginning of the `run0` script (i.e. set the value of `nummax`)
- Copy start files `start.nc` `startfi.nc` over and rename them `start0.nc` `startfi0.nc`.
- Run script `run0`

`run0` runs a series of simulations that generate the indexed output files (e.g. `start1`, `startfi1`, `diagfi1`, etc.) including files `lrun1`, `lrun2`, etc. containing the redirection of the display and the information about the run.

NOTE: to restart a series of simulations after a first series (for example, starting from `start5` and `startfi5`), just write the index of the initial files (e.g. 5) in the file named `num.run`. If `num.run` exists, the model will start from the index written in `num.run`. If not it will start from `start0` and `startfi0`.

NOTE: A script is available for performing annual runs with 12 seasons at 30° solar longitude as it is in the database (script `run_mcd`, also found in directory `deftank`). This script functions with script `run0`. Just set the number of simulations to 1 in `run0`. Then copy `run.def` into `run.def.ref` and set `nday` to 9999 in this file. To start from `startN.c`, edit the file `run_mcd` and comment (with a #) the N months already created and describe N in `num.run`. Then run `run_mcd`.

4.10 Creating and modifying initial states

4.10.1 Using program “newstart”

Several model parameters (for example, the dust optical depth) are stored in the initial states (NetCDF files `start.nc` and `startfi.nc`). To change these parameters, or to generally change the model resolution, use program **newstart**.

This program is also used to create an initial state. In practice, we usually reuse an old initial state, and modify it using **newstart**.

Like the GCM, program **newstart** must be compiled (using the `makelmdz_fcm` or `makegcm` script) at the required grid resolution. For example:

```
makelmdz_fcm -arch local -d 64x48x25 -p mars newstart
```

Then run the `newstart` program in a directory containing the `start` and `def` file to be used:

```
newstart.e
```

The program then gives you two options:

```
A partir de quoi souhaitez vous creer vos etats initiaux ?
  0 - d un fichier start_archive
  1 - d un fichier start
```

- - Option “1” allows you to read and modify the information needed to create a new initial state from the files `start.nc`, `startfi.nc`
- - Option “0” allows you to read and modify the information needed to create a new initial state from file `start_archive.nc` (whatever the `start_archive.nc` grid resolution is).

If you use tracers, make sure that they are taken into account in your start files (either `start` or `start_archive`).

Then answer to the various questions in the scroll menu. These questions allow you to modify the initial state for the following parameters.

```
First set of questions:
Modifications of variables in tab_cntrl:
~~~~~
day_ini : Jour initial (=0 a Ls=0)
z0 : surface roughness (m)
emin_turb : energie minimale
lmixmin : longueur de melange
emissiv : Emissivite du sol martien
emisice : Emissivite des calottes
albedice : Albedo des calotte
iceradius : mean scat radius of CO2 snow
dtemisice : time scale for snow,
. ' metamorphism
    tauvis : profondeur optique visible,
. ' moyenne
obliquit : planet obliquity (deg)
peri_day : perihelion date (sol since Ls=0)
periheli : min. sun-mars dist (Mkm)
aphelie : max. sun-mars dist (Mkm)
```

```
Second set of questions :
flat : no topography ("aquaplanet")
bilball : albedo, inertie thermique uniforme
coldspole : sous sol froid et haut albedo au pole sud
```

```

q=0 : traceurs a zero
ini_q : traceurs initialises pour la chimie
ini_q-H2O : idem, sauf le dernier traceur (H2O)
ini_q-iceH2O : idem, sauf ice et H2O
watercapn : H2O ice sur la calotte permanente nord
watercaps : H2O ice sur la calotte permanente sud
wetstart : start with a wet atmosphere
iqset : give a specific value to tracer iq
isotherm : Temperatures isothermes et vents nuls
co2ice=0 : elimination des calottes polaires de CO2
ptot : pression totale

```

Program **newstart.e** creates files `restart.nc` and `restartfi.nc` that you generally need to rename (for instance rename them in `start0.nc` and `startfi0.nc` if you want to use `run0` or `run_mcd`, starting with season 0; rename them `start.nc` and `startfi.nc` if you just want to perform one run with `gcm.e`).

4.10.2 Creating the initial `start_archive.nc` file

Archive file `start_archive.nc` is created from files `start.nc` and `startfi.nc` by program **start2archive**. Program **start2archive** compiles to the same grid resolution as the `start.nc` and `startfi.nc` grid resolution. For example:

```
makelmd_fcm -arch local -d 64x48x25 -p mars start2archive
```

Then run `start2archive.e`

You now have a `start_archive.nc` file for one season that you can use with `newstart`. If you want to gather other states obtained at other times of year, rerun `start2archive.e` with the `start.nc` and `startfi.nc` corresponding to these. These additional initial states will automatically be added to the `start_archive.nc` file present in the directory.

4.10.3 Changing the horizontal or vertical grid resolution

To run at a different grid resolution than available initial conditions files, one needs to use tools **newstart** and **start2archive**

For example, to create initial states at grid resolution $32 \times 24 \times 25$ from NetCDF files `start` and `startfi` at grid resolution $64 \times 48 \times 32$:

- Create file `start_archive.nc` with **start2archive.e** compiled at grid resolution $64 \times 48 \times 25$ using **old file** `z2sig.def` **used previously**
- Create files `newstart.nc` and `newstartfi.nc` with **newstart.e** compiled at grid resolution $32 \times 24 \times 25$, using **new file** `z2sig.def`

If you want to create starts files with tracers for 49 layers using a `start_archive.nc` obtained for 32 layers, do not forget to use the `ini_q` option in `newstart` in order to correctly initialize tracers value for layer 33 to layer 49. You just have to answer yes to the question on thermosphere initialization if you want to initialize the thermosphere part only ($l=33$ to $l=49$), and no if you want to initialize tracers for all layers ($l=0$ to $l=49$).

Chapter 5

Program organization and compilation script

Currently the model is split between two directories: **LMDZ.COMMON** and **LMDZ.MARS**. The reason for this separation is that all that concerns the dynamics (and which is common to all planets) is in **LMDZ.COMMON** whereas all that concerns the Martian physics is in **LMDZ.MARS**. For historical reasons, **LMDZ.MARS** still includes a dynamical core, but it is strongly recommended to use the **LMDZ.COMMON** (mandatory if you want to run in parallel).

Here is a brief description of the contents of directory **LMDZ.COMMON**:

```
libf/  model FORTRAN Source code (.F or .F90)
        and include files (.h) organised in sub-directories
        (dynamics (dyn3d), filters (filtrez), interface with
        physical packages (dynphy_lonlat), ...)
```

```
arch/  set of architecture files used by the makelmdz_fcm script
```

```
makelmdz_fcm  script to use to compile the GCM and related
              utilities
```

```
bin/  directory where executables generated by the makelmdz_fcm
       script will be generated
```

```
ioipsl/ directory with scripts usefull to download and build the
        IOIPSL library (not needed by the Martian GCM).
```

And a brief description of the **LMDZ.MARS** directory contents:

```
libf/  Model FORTRAN Source code (.F or .F90)
        and include files (.h) organised in sub-directories
        (physics (phymars), dynamics (dyn3d), filters (filtrez)...)

deftank/ A collection of examples of parameter files required
         to run the GCM (run.def, callphys.def, ...)
```

```
util/  A set of programs useful for post-processing GCM outputs.
```

```
makegcm (Obsolete) script that should be used to compile the GCM as well
         as related utilities (newstart, start2archive, testphysld)
```

5.1 Organization of the model source files

The model source files are stored in various sub directories in directory **libf**. These sub-directories correspond to the different parts of the model:

grid: mainly made up of "dimensions.h" file, which contains the parameters that define the model grid, i.e. the number of points in longitude (IIM), latitude (JJM) and altitude (LLM), as well as the number of tracers (NQMX).

dyn3d: contains the dynamical subroutines.

biblio: contains some generic subroutines not specifically related to physics or dynamics but used by either or both.

phymars: contains the Martian physics routines.

filtrez: contains the longitudinal filter sources applied in the upper latitudes, where the Courant-Friedrich-Levy stability criterion is violated.

aeronomars: contains the Martian chemistry and thermosphere routines.

5.2 Programming

The model is written in **Fortran-77** and **Fortran-90**.

- The program sources are written in "file.F" or "file.F90" files. The extension .F is the standard extension for fixed-form Fortran and the extension .F90 is for free-form Fortran. These files must be preprocessed (by a **C preprocessor** such as (cpp)) before compilation (this behaviour is, for most compilers, implicitly obtained but using a capital F in the extension of the file names).
- Constants are placed in COMMON declarations, located in the common "include" files "file.h"
- In general, variables are passed from subroutine to subroutine as arguments (and never as COMMON blocks).
- In some parts of the code, for "historical" reasons, the following rule is sometimes used: in the subroutine, the variables (ex: name) passed as an argument by the calling program are given the prefix p (ex: pname) while the local variables are given the prefix z (ex: zname). As a result, several variables change their prefix (and thus their name) when passing from a calling subroutine to a called subroutine.

5.3 Model organization

Figure 5.1 describes the main subroutines called by physiq.F.

5.4 Compiling the model

Technically, the model is compiled using the Unix utility `make`. The `make1mdz_fcm` utility script recreates the `makefile` file when necessary, for example, when a source file has been added or removed since the last compilation.

None of this is visible to the user. To (re-)compile the model just run the command

```
make1mdz_fcm
```

with adequate options (e.g. `make1mdz_fcm -arch loacl -d 62x48x32 -p mars gcm`), as described in section 4.4.




Figure 5.1: Organigram of subroutine function physiq.F

Help manual for the makelmdz_fcm script

Use the "-h" option to learn about possible options:

```
make1mdz_fcm -h
```

Chapter 6

Input/Output

6.1 NetCDF format

GCM input/output data are written in **NetCDF** format (Network Common Data Form). NetCDF is an interface used to store and access geophysical data, and a library that provides an implementation of this interface. The NetCDF library also defines a machine-independent format for representing scientific data. Together, the interface, library and format support the creation, access and sharing of scientific data. NetCDF was developed at the Unidata Program Center in Boulder, Colorado. The freely available source can be obtained from the Unidata website.

<http://www.unidata.ucar.edu/software/netcdf>

A data set in NetCDF format is a single file, as it is self-descriptive.

6.1.1 NetCDF file editor: `ncdump`

The editor is included in the NetCDF library. By default it generates an ASCII representation as standard output from the NetCDF file specified at the input.

Main commands for `ncdump`

`ncdump diagfi.nc`

dump contents of NetCDF file `diagfi.nc` to standard output (i.e. the screen).

`ncdump -c diagfi.nc`

Displays the **coordinate** variable values (variables which are also dimensions), as well as the declarations, variables and attribute values. The values of the non-coordinate variable data are not displayed at the output.

`ncdump -h diagfi.nc`

Shows only the informative header of the file, which is the declaration of the dimensions, variables and attributes, but not the values of these variables. The output is identical to that in option **-c** except for the fact that the coordinated variable values are not included.

`ncdump -v var1,...,varn diagfi.nc`

The output includes the specific variable values, as well as all the dimensions, variables and attributes. More than one variable can be specified in the list following this option. The list must be a simple argument for the command, and must not contain any spaces. If no variable is specified, the command displays all the values of the variables in the file by default.




Figure 6.1: Example of temperature data at a given time using GrADS visualization

6.1.2 Graphic visualization of the NetCDF files using GrAds

GrAdS (The Grid Analysis and Display System) is a graphic software developed by Brian Doty at the "Center for Ocean-Land-Atmosphere (COLA)".

One of its functions is to enable data stored in NetCDF format to be visualized directly. In figure 6.1 for example, we can see the GrADS visualization of the temperature data at a given moment. However, unlike NetCDF, GrADS only recognizes files where all the variables are stored on the same horizontal grid. These variables can be in 1, 2, 3 or 4 dimensions (X,Y,Z and t).

GrADS can also be obtained on the WWW.

<http://grads.iges.org/grads/>

6.2 Input and parameter files

The (3D version of the) GCM requires the input of two initialization files (in NetCDF format):

-**start.nc** contains the initial states of the dynamical variables.

-**startfi.nc** contains the initial states of the physical variables.

Note that collections of initial states can be retrieved at:

<http://www.lmd.jussieu.fr/~forget/datagcm/Starts>

Extracting **start.nc** and **startfi.nc** from these archived requires using program **newstart**, as described in section 4.10.

To run, the GCM also requires the four following parameter files (ascii text files):

-**run.def** the parameters of the dynamical part of the program, and the temporal integration of the model.

-**callphys.def** the parameters for calling the physical part.

-**traceur.def** the names of the tracer to use.

-**z2sig.def** the vertical distribution of the atmospheric layers.

Examples of these parameter files can be found in the **LMDZ.MARS/def/tank** directory.

6.2.1 run.def

A typical `run.def` file is given as an example below. The choice of variables to be set is simple (e.g. `nday` number of modeled days to run), while the others do not need to be changed for normal use.

The format of the `run.def` file is quite straightforward (and flexible): values given to parameters must be given as:

```
parameter = value
```

Any blank line or line beginning with symbol `#` is a comment, and instruction lines may be written in any order. Moreover, not specifying a parameter/value set (e.g. deleting it or commenting it out) means you want the GCM to use a default built-in value. Additionally, one may use a specific keyword **INCLUDEDEF** to specify another (text) file in which to also read values of parameters; e.g.:

```
INCLUDEDEF=callphys.def
```

Here are some details about some of the parameters which may be set in `run.def`:

- **day_step**, the number of dynamical steps per day to use for the time integration. This needs to be large enough for the model to remain stable (this is related to the CFL stability criterion which essentially depends on the horizontal resolution of the model). On Mars, in theory, the GCM can run with `day_step=480` using the 64×48 grid, but model stability improves when this number is higher: `day_step=960` is recommended when using the 64×48 grid. According to the CFL criterion, `day_step` should vary in proportion with the resolution: for example `day_step=480` using the 32×24 horizontal resolution. Note that `day_step` must also be divisible by `iperiod`.
- **tetagdiv**, **tetagrot**, **tetatemp** control the dissipation intensity. It is better to limit the dissipation intensity (`tetagdiv`, `tetagrot`, `tetatemp` should not be too low). However the model diverges if `tetagdiv`, `tetagrot`, `tetatemp` are too high, especially if there is a lot of dust in the atmosphere.
Example used with `nitergdiv=1` and `nitergrot=niterh=2` :
 - using the 32×24 grid `tetagdiv=6000 s` ; `tetagrot=tetatemp=30000 s`
 - using the 64×48 grid: `tetagdiv=3000 s` ; `tetagrot=tetatemp=9000 s`
- **idissip** is the time step used for the dissipation: dissipation is computed and added every `idissip` dynamical time step. If `idissip` is too short, the model waste time in these calculations. But if `idissip` is too long, the dissipation will not be parametrized correctly and the model will be more likely to diverge. A check must be made, so that: `idissip < tetagdiv*daystep/88775` (same rule for `tetagrot` and `tetatemp`). This is tested automatically during the run.
- **iphysiq** is the time step used for the physics: physical tendencies are computed every `iphysiq` dynamical time step. In practice, we usually set the physical time step to be of the order of half an hour. We thus generally set `iphysiq=day_step/48`

Example of run.def file:

```
#
#-----
#GCM run control parameters:
#-----

# planet type
planet_type=mars

# Number of days to run model for
```

```

    nday=9999

# Number of dynamical steps per day (must be a multiple of iperiod)
day_step = 960

# Apply a Matsuno step every iperiod dynamical step
iperiod=5

# Control output information in the dynamics every iconser dynamical steps
iconser=120

# Apply dissipation every idissip dynamical steps
idissip=1

# dissipation operator to use (star or non-star)
lstaridis=.true.

# use hybrid vertical coordinate (else will use sigma levels)
hybrid=.true.

# iterate lateral dissipation operator gradiv nitergdiv times
nitergdiv=1

# iterate lateral dissipation operator nxgradrot nitergrot times
nitergrot=2

# iterate lateral dissipation operator divgrad niterh times
niterh=2

# time scale (s) for shortest wavelengths for u,v (gradiv)
tetagdiv= 2500.

# time scale (s) for shortest wavelengths for u,v (nxgradrot)
tetagrot=5000.

# time scale (s) for shortest wavelengths for h (divgrad)
tetatemp=5000.

# coefficient for gamdissip
coefdis=0.

# time marching scheme (Matsuno if purmats is true, else Matsuno-Leapfrog)
purmats=.false.

# run with (true) or without (false) physics
physic=.true.

# call physics every iphysiq dynamical steps
iphysiq=10

# Use a regular grid
grireg=.true.

# Output in diagfi file every ecritphy dynamical steps
ecritphy=240

# longitude (degrees) of zoom center
clon=63.

# latitude (degrees) of zoom center
clat=0.

# enhancement factor of zoom, along longitudes
grossismx=1.

# enhancement factor of zoom, along latitudes
grossismy=1.

```



```

# Use an hyperbolic function f(y) if .true., else use a sine
fxyhypb=.false.

# extention along longitudes of zoom region (fraction of global domain)
dzoomx= 0.

# extention along latitudes of zoom region (fraction of global domain)
dzoomy=0.

# zoom stiffness along longitudes
taux=2.

# zoom stiffness along latitudes
tauy=2.

# Fonction f(y) as y = Sin(latitude) if = .true. , else y = latitude
ysinus= .false.

# Use a sponge layer
callsponge = .true.

# Sponge layer extends over topmost nsponge layers
nsponge = 3

# Sponge: mode0(u=v=0), mode1(u=umoy,v=0), mode2(u=umoy,v=vmoy)
mode_sponge= 3

# Sponge layer time scale (s): tetasponge
tetasponge = 30000

# some definitions for the physics, in file 'callphys.def'
INCLUDEDEF=callphys.def

```

6.2.2 callphys.def

The `callphys.def` file (along the same format as the `run.def` file) contains parameter/value sets for the physics.

Example of callphys.def file:

```

##General options
##~~~~~
#Run with or without tracer transport ?
tracer=.true.

#Directory where external input files are:
datadir=./datafile

#Diurnal cycle ? if diurnal=False, diurnal averaged solar heating
diurnal=.true.

#Seasonal cycle ? if season=False, Ls stays constant, to value set in "start"
season = .true.

#write some more output on the screen ?
lwrite = .false.

#Save statistics in file "stats.nc" ?
callstats =.true.

#Save EOF profiles in file "profiles" for Climate Database?
calleofdump = .false.

```

```

## Dust scenario. Used if the dust is prescribed (i.e. if tracer=F or active=F)
## ~~~~~
# =1 Dust opt.deph read in startfi; =2 Viking scenario; =3 MGS scenario,
# =4 Mars Year 24 from TES assimilation (old version of MY24; dust_tes.nc file)
# =6 "cold" (low dust) scenario ; =7 "warm" (high dust) scenario
# =8 "clim" scenario
# =24 Mars Year 24 ; =25 Mars Year 25 (year with a global dust storm) ; ...
# =30 Mars Year 30
iaervar = 32
# Dust opacity at 610 Pa (when constant, i.e. for the iaervar=1 case)
tauvis=0.1
# Dust vertical distribution:
# (=0: old distrib. (Pollack90), =1: top set by "topdustref",
# =2: Viking scenario; =3 MGS scenario)
iddist = 3
# Dust top altitude (km). (Matters only if iddist=1)
topdustref = 55.

## Physical Parameterizations :
## ~~~~~
# call radiative transfer ?
callrad = .true.
# call NLTE radiative schemes ? matters only if callrad=T
callnlte = .true.
# NLTE 15um scheme to use.
# 0-> Old scheme, static oxygen
# 1-> Old scheme, dynamic oxygen
# 2-> New scheme
nltemodel = 2
# call CO2 NIR absorption ? matters only if callrad=T
callnirco2 = .true.
# NIR NLTE correction ? matters only if callnirco2=T
nircorr=1
# call turbulent vertical diffusion ?
calldifv = .true.
# call convective adjustment ?
calladj = .true.
# Thermals
calltherm = .true.
callrichsl = .true.
# call CO2 condensation ?
callcond =.true.
# call thermal conduction in the soil ?
callsoil = .true.
# call Lott's gravity wave/subgrid topography scheme ?
calllott = .true.
# Impose polar cap surface albedos as observed by TES?
TESicealbedo = .true.
## Coefficient for Northern cap albedoes
TESice_Ncoef=1.6
## Coefficient for Southern cap albedoes
TESice_Scoef=1.6

## Radiative transfer options :
## ~~~~~
# the rad.transfer is computed every "iradia" physical timestep
iradia = 1
# Output of the exchange coefficient matrix ? for diagnostic only
callg2d = .false.
# Rayleigh scattering : (should be .false. for now)
rayleigh = .false.
#number of radiatively active tracers
naerkind = 2

## Tracer (dust water, ice and/or chemical species) options (used if tracer=T):
## ~~~~~

```

```

# DUST: Transported dust ? (if >0, use 'dustbin' dust bins)
dustbin = 2
# DUST: Radiatively active dust ? (matters if dustbin>0)
active = .true.
# DUST: use mass and number mixing ratios to predict dust size ?
doubleq = .true.
# DUST: lifted by GCM surface winds ?
lifting = .true.
# DUST: lifted by dust devils ?
callddevil = .false.
# DUST: Scavenging by H2O/CO2 snowfall ?
scavenging = .true.
# DUST/WATERICE: Gravitational sedimentation ?
sedimentation = .true.
# WATERICE: Radiatively active transported atmospheric water ice ?
activice = .true.
# WATER: Compute water cycle
water = .true.
# WATER: Microphysical scheme for water-ice clouds?
microphys = .true.
# WATER: parameter contact
mteta = 0.95
# WATER: current permanent caps at both poles. True IS RECOMMENDED
#       (with .true., North cap is a source of water and South pole
#       is a cold trap)
caps = .true.
# WATER: Water ice albedo ?
albedo_h2o_ice = 0.35
# WATER: Water ice thermal inertia
inert_h2o_ice = 800
# WATER: Frost thickness threshold for albedo
frost_albedo_threshold = 0.005
nuice_sed=0.1
icedryness=1

# PHOTOCHEMISTRY: include chemical species
photochem = .true.

## Thermospheric options (relevant if tracer=T) :
##~~~~~
# call thermosphere ?
callthermos = .true.
# WATER: included without cycle (only if water=.false.)
thermoswater = .false.
# call thermal conduction ? (only if callthermos=.true.)
callconduct = .true.
# call molecular viscosity ? (only if callthermos=.true.)
callmolvis = .true.
# call molecular diffusion ? (only if callthermos=.true.)
callmoldiff = .true.
# call thermospheric photochemistry ? (only if callthermos=.true.)
thermochem = .true.
# call EUV heating ? (only if callthermos=.true.)
calleuv=.true.
#Method to include solar variability?
#0-> Old method (fixed EUV input) 1-> Variability with E10.7 as observed
solvarmod=1
# date for solar flux calculation: (1985 < date < 2002)
## (Solar min=1996.4 ave=1993.4 max=1990.6) ; Only used if solvarmod=0
solarcondate = 1993.4
#Solar variability as observed for MY? (must bebetween MY23 and MY30)
# (only matters if solvarmod=1)
solvaryear=32
# value for the UV heating efficiency
##(experimental values between 0.19 and 0.23, lower values may
## be used to compensate for low 15 um cooling)
euveff = 0.21

```

6.2.3 traceur.def

Tracers in input (`start.nc` and `startfi.nc`) and output files (`restart.nc` and `restartfi.nc`) are stored using individual tracer names (e.g. `co2` for CO2 gas, `h2o_vap` for water vapour, `h2o_ice` for water ice, ...).

The first line of the `traceur.def` file (an ASCII file) must contain the number of tracers to load and use (this number should be the same as given to the `-t` option of the `makegcm` script when the GCM was compiled), followed by the tracer names (one per line). Note that if the corresponding tracers are not found in input files `start.nc` and `startfi.nc`, then the tracer is initialized to zero.

Example of a traceur.def file: (with water vapour and ice tracers)

```
2
h2o_ice
h2o_vap
```

6.2.4 z2sig.def

The `z2sig.def` file contains the pseudo-altitudes (in km) at which the user wants to set the vertical levels.

Note that levels should be unevenly spread, with a higher resolution near the surface in order to capture the rapid variations of variables there. It is recommended to use the altitude levels as set in the `z2sig.def` file provided in the `deftank` directory.

Example of z2sig.def file (this version for 49 layers between 0 and 300 km):

```
10.00000      H: atmospheric scale height (km) (used as a reference only)
0.0020736
0.015735
0.0600151
0.163344
0.36358
0.708007
1.25334
2.06571
3.2
4.7
6.6
8.9
11.6
14.7
18.2
22.1
26.4
31.1
36.2
41.7
47.6
53.9
60.55
67.5
74.5
81.5
88.5
95.5
```

102.5
109.5
116.5
123.5
130.5
137.5
144.5
151.5
158.5
165.5
172.5
179.5
186.5
193.5
200.5
207.5
214.5
221.5
228.5
235.5
242.5

6.2.5 Initialization files: start and startfi

Files `start.nc` and `startfi.nc`, like all the NetCDF files of the GCM, are constructed on the same model (see NetCDF file composition, figure 6.2). They contain:

- a header with a “control” variable followed by a series of variables defining the (physical and dynamical) grids
- a series of non temporal variables that give information about surface conditions on the planet.
- a “time” variable giving the values of the different instants at which the temporal variables are stored (a single time value (t=0) for start, as it describes the dynamical initial states, and no time values for startfi, as it describes only a physical state).

To visualize the contents of a `start.nc` file using the `ncdump` command:

```
ncdump -h start.nc
```

```
netcdf start {  
  dimensions:  
    index = 100 ;  
    rlonu = 33 ;  
    latitude = 25 ;  
    longitude = 33 ;  
    rlatv = 24 ;  
    altitude = 18 ;  
    interlayer = 19 ;  
    Time = UNLIMITED ; // (1 currently)  
  variables:  
    float controle(index) ;  
      controle:title = "Parametres de controle" ;  
    float rlonu(rlonu) ;  
      rlonu:title = "Longitudes des points U" ;  
    float rlatu(latitude) ;  
      rlatu:title = "Latitudes des points U" ;  
    float rlonv(longitude) ;  
      rlonv:title = "Longitudes des points V" ;  
    float rlatv(rlatv) ;
```




Figure 6.2: Organization of NetCDF files

```

      rlatv:title = "Latitudes des points V" ;
float ap(interlayer) ;
      ap:title = "Coef A: hybrid pressure levels" ;
float bp(interlayer) ;
      bp:title = "Coef B: hybrid sigma levels" ;
float aps(altitude) ;
      aps:title = "Coef AS: hybrid pressure at midlayers" ;
float bps(altitude) ;
      bps:title = "Coef BS: hybrid sigma at midlayers" ;
float presnivs(altitude) ;
float latitude(latitude) ;
      latitude:units = "degrees_north" ;
      latitude:long_name = "North latitude" ;
float longitude(longitude) ;
      longitude:long_name = "East longitude" ;
      longitude:units = "degrees_east" ;
float altitude(altitude) ;
      altitude:long_name = "pseudo-alt" ;
      altitude:units = "km" ;
      altitude:positive = "up" ;

```

```

float cu(latitude, rlonu) ;
    cu:title = "Coefficient de passage pour U" ;
float cv(rlatv, longitude) ;
    cv:title = "Coefficient de passage pour V" ;
float aire(latitude, longitude) ;
    aire:title = "Aires de chaque maille" ;
float phisinit(latitude, longitude) ;
    phisinit:title = "Geopotentiel au sol" ;
float Time(Time) ;
    Time:title = "Temps de simulation" ;
    Time:units = "days since 1-01-01 00:00:00" ;
float ucov(Time, altitude, latitude, rlonu) ;
    ucov:title = "Vitesse U" ;
float vcov(Time, altitude, rlatv, longitude) ;
    vcov:title = "Vitesse V" ;
float teta(Time, altitude, latitude, longitude) ;
    teta:title = "Temperature" ;
float h2o_ice(Time, altitude, latitude, longitude) ;
    h2o_ice:title = "Traceur h2o_ice" ;
float h2o_vap(Time, altitude, latitude, longitude) ;
    h2o_vap:title = "Traceur h2o_vap" ;
float masse(Time, altitude, latitude, longitude) ;
    masse:title = "C est quoi ?" ;
float ps(Time, latitude, longitude) ;
    ps:title = "Pression au sol" ;

// global attributes:
    :title = "Dynamic start file" ;
}

```

List of contents of a `startfi.nc` file:

ncdump -h startfi.nc

```

netcdf startfi {
dimensions:
    index = 100 ;
    physical_points = 738 ;
    subsurface_layers = 18 ;
    nlayer_plus_1 = 19 ;
    number_of_advected_fields = 3 ;
variables:
    float controle(index) ;
        controle:title = "Control parameters" ;
    float soildepth(subsurface_layers) ;
        soildepth:title = "Soil mid-layer depth" ;
    float longitude(physical_points) ;
        longitude:title = "Longitudes of physics grid" ;
    float latitude(physical_points) ;
        latitude:title = "Latitudes of physics grid" ;
    float area(physical_points) ;
        area:title = "Mesh area" ;
    float phisfi(physical_points) ;
        phisfi:title = "Geopotential at the surface" ;
    float albedodat(physical_points) ;
        albedodat:title = "Albedo of bare ground" ;
    float ZMEA(physical_points) ;
        ZMEA:title = "Relief: mean relief" ;
    float ZSTD(physical_points) ;
        ZSTD:title = "Relief: standard deviation" ;
    float ZSIG(physical_points) ;
        ZSIG:title = "Relief: sigma parameter" ;
    float ZGAM(physical_points) ;
        ZGAM:title = "Relief: gamma parameter" ;
    float ZTHE(physical_points) ;
        ZTHE:title = "Relief: theta parameter" ;
    float co2ice(physical_points) ;

```

```

        co2_ice:title = "CO2 ice cover" ;
float inertiedat(subsurface_layers, physical_points) ;
        inertiedat:title = "Soil thermal inertia" ;
float tsurf(physical_points) ;
        tsurf:title = "Surface temperature" ;
float tsoil(subsurface_layers, physical_points) ;
        tsoil:title = "Soil temperature" ;
float emis(physical_points) ;
        emis:title = "Surface emissivity" ;
float q2(nlayer_plus_1, physical_points) ;
        q2:title = "pbl wind variance" ;
float h2o_ice(physical_points) ;
        h2o_ice:title = "tracer on surface" ;

// global attributes:
        :title = "Physics start file" ;
}

```

Physical and dynamical headers

There are two types of headers: one for the physical headers, and one for the dynamical headers. The headers always begin with a “control” variable (described below), that is allocated differently in the physical and dynamical parts. The other variables in the header concern the (physical and dynamical) grids. They are the following:

the horizontal coordinates

- **rlonu, rlatu, rlonv, rlatv** for the dynamical part,
- **lati, long** for the physical part,

the coefficients for passing from the physical grid to the dynamical grid

- **cu,cv** only in the dynamical header

and finally, the grid box areas

- **aire** for the dynamical part,
- **area** for the physical part.

Surface conditions

The surface conditions are mostly given in the physical NetCDF files by variables:

- **phisfi** for the initial state of surface geopotential,
- **albedodat** for the bare ground albedo,
- **inertiedat** for the surface thermal inertia,
- **zmea, zstd, zsig, zgam** and **zthe** for the subgrid scale topography.

For the dynamics:

- **physinit** for the initial state of surface geopotential

Remark: variables **phisfi** and **physinit** contain the same information (surface geopotential), but **phisfi** gives the geopotential values on the physical grid, while **physinit** give the values on the dynamical grid.

Physical and dynamical state variables

To save disk space, the initialization files store the variables used by the model, rather than the “natural” variables.

For the dynamics:

- **ucov** and **vcov** the covariant winds
 These variables are linked to the “natural” winds by

$$ucov = cu * u \text{ and } vcov = cv * v$$
- **teta** the potential temperature,
 or more precisely, the potential enthalpy linked to temperature **T** by $\theta = T \left(\frac{P}{P_{ref}} \right)^{-K}$
- the tracers,
- **ps** surface pressure.
- **masse** the atmosphere mass in each grid box.

“Vectorial” variables **ucov** and **vcov** are stored on “staggered” grids u and v respectively (in the dynamics) (see section 2.2).

Scalar variables **h**, **q** (tracers), **ps**, **masse** are stored on the “scalar” grid of the dynamical part.

For the physics:

- **co2ice** surface dry ice,
- **tsurf** surface temperature,
- **tsoil** temperatures at different layers under the surface,
- **emis** surface emissivity,
- **q2** wind variance,
 or more precisely, the square root of the turbulent kinetic energy.
- the surface “tracer” budget (kg.m^{-2}),

All these variables are stored on the “physical” grid (see section 2.2).

The “control” array

Both physical and dynamical headers of the GCM NetCDF files start with a **controle** variable. This variable is an array of 100 reals (the vector called `tab_cntrl` in the program), which contains the program control parameters. Parameters differ between the physical and dynamical sections, and examples of both are listed below. The contents of table `tab_cntrl` can also be checked with the command `ncdump -ff -v controle`.

The “control” array in the header of a dynamical NetCDF file: start

```

tab_cntrl(1) = FLOAT(iim) ! number of nodes along longitude
tab_cntrl(2) = FLOAT(jjm) ! number of nodes along latitude
tab_cntrl(3) = FLOAT(llm) ! number of atmospheric layers
tab_cntrl(4) = FLOAT(idayref) ! initial day
tab_cntrl(5) = rad    ! radius of the planet
tab_cntrl(6) = omeg   ! rotation of the planet (rad/s)
tab_cntrl(7) = g      ! gravity (m/s2) ~3.72 for Mars
tab_cntrl(8) = cpp
tab_cntrl(9) = kappa  ! = r/cp
tab_cntrl(10) = daysec ! lenght of a sol (s) ~88775
tab_cntrl(11) = dtvr  ! dynamical time step (s)
tab_cntrl(12) = etot0 ! total energy
tab_cntrl(13) = ptot0 ! total pressure
tab_cntrl(14) = ztot0 ! total enstrophy

```

```

tab_cntrl(15) = stot0 ! total enthalpy
tab_cntrl(16) = ang0 ! total angular momentum
tab_cntrl(17) = pa
tab_cntrl(18) = preff ! reference pressure (Pa)
tab_cntrl(19) = clon ! longitude of center of zoom
tab_cntrl(20) = clat ! latitude of center of zoom
tab_cntrl(21) = grossismx ! zooming factor, along longitude
tab_cntrl(22) = grossismy ! zooming factor, along latitude

tab_cntrl(24) = dzoomx ! extention (in longitude) of zoom
tab_cntrl(25) = dzoomy ! extention (in latitude) of zoom

tab_cntrl(27) = taux ! stiffness factor of zoom in longitude
tab_cntrl(28) = tauy ! stiffness factor of zoom in latitude

```

The "controle" array in the header of a physical NetCDF file: startfi.nc

```

c Informations on the physics grid
  tab_cntrl(1) = float(ngridmx) ! number of nodes on physics grid
  tab_cntrl(2) = float(nlayermx) ! number of atmospheric layers
  tab_cntrl(3) = day_ini + int(time) ! initial day
  tab_cntrl(4) = time -int(time) ! initiale time of day

c Informations about Mars, used by dynamics and physics
  tab_cntrl(5) = rad ! radius of Mars (m) ~3397200
  tab_cntrl(6) = omeg ! rotation rate (rad.s-1)
  tab_cntrl(7) = g ! gravity (m.s-2) ~3.72
  tab_cntrl(8) = mugaz ! Molar mass of the atmosphere (g.mol-1) ~43.49
  tab_cntrl(9) = rcp ! = r/cp ~0.256793 (=kappa dans dynamique)
  tab_cntrl(10) = daysec ! length of a sol (s) ~88775

  tab_cntrl(11) = phystep ! time step in the physics
  tab_cntrl(12) = 0.
  tab_cntrl(13) = 0.

c Informations about Mars, only for physics
  tab_cntrl(14) = year_day ! length of year (sols) ~668.6
  tab_cntrl(15) = periheli ! min. Sun-Mars distance (Mkm) ~206.66
  tab_cntrl(16) = aphelie ! max. Sun-Mars distance (Mkm) ~249.22
  tab_cntrl(17) = peri_day ! date of perihelion (sols since N. spring)
  tab_cntrl(18) = obliquit ! Obliquity of the planet (deg) ~23.98

c Boundary layer and turbulence
  tab_cntrl(19) = z0 ! surface roughness (m) ~0.01
  tab_cntrl(20) = lmixmin ! mixing length ~100
  tab_cntrl(21) = emin_turb ! minimal energy ~1.e-8

c Optical properties of polar caps and ground emissivity
  tab_cntrl(22) = albedice(1) ! Albedo of northern cap ~0.5
  tab_cntrl(23) = albedice(2) ! Albedo of southern cap ~0.5
  tab_cntrl(24) = emisice(1) ! Emissivity of northern cap ~0.95
  tab_cntrl(25) = emisice(2) ! Emissivity of southern cap ~0.95
  tab_cntrl(26) = emissiv ! Emissivity of martian soil ~.95
  tab_cntrl(31) = iceradius(1) ! mean scat radius of CO2 snow (north)
  tab_cntrl(32) = iceradius(2) ! mean scat radius of CO2 snow (south)
  tab_cntrl(33) = dtemisice(1) ! time scale for snow metamorphism (north)
  tab_cntrl(34) = dtemisice(2) ! time scale for snow metamorphism (south)

c dust aerosol properties
  tab_cntrl(27) = tauvis ! mean visible optical depth

  tab_cntrl(28) = 0.
  tab_cntrl(29) = 0.
  tab_cntrl(30) = 0.

! Soil properties:
  tab_cntrl(35) = volcapa ! soil volumetric heat capacity

```

6.3 Output files

6.3.1 NetCDF restart files - restart.nc and restartfi.nc

These files are of the exact same format as `start.nc` and `startfi.nc`

6.3.2 NetCDF file - diagfi.nc

NetCDF file `diagfi.nc` stores the instantaneous physical variables throughout the simulation at regular intervals (set by the value of parameter `ecritphy` in parameter file `run.def`; note that `ecritphy` should be a multiple of `iphysiq` as well as a divisor of `day_step`).

Any variable from any sub-routine of the physics can be stored by calling subroutine `writediagfi`. Moreover, one may add a `diagfi.def` file containing only the names of variables to output (one per line) in the directory where the GCM is run, in order to have only those listed outputted in the `diagfi.nc` file.

Illustrative example of the contents of a `diagfi.nc` file (using `ncdump`):
ncdump -h diagfi.nc

```
netcdf diagfi {
dimensions:
    Time = UNLIMITED ; // (12 currently)
    index = 100 ;
    rlonu = 65 ;
    latitude = 49 ;
    longitude = 65 ;
    rlatv = 48 ;
    interlayer = 26 ;
    altitude = 25 ;
    subsurface_layers = 18 ;
variables:
    float Time(Time) ;
        Time:long_name = "Time" ;
        Time:units = "days since 0000-00-0 00:00:00" ;
    float controle(index) ;
        controle:title = "Control parameters" ;
    float rlonu(rlonu) ;
        rlonu:title = "Longitudes at u nodes" ;
    float latitude(latitude) ;
        latitude:units = "degrees_north" ;
        latitude:long_name = "North latitude" ;
    float longitude(longitude) ;
        longitude:long_name = "East longitude" ;
        longitude:units = "degrees_east" ;
    float altitude(altitude) ;
        altitude:long_name = "pseudo-alt" ;
        altitude:units = "km" ;
        altitude:positive = "up" ;
    float rlatv(rlatv) ;
        rlatv:title = "Latitudes at v nodes" ;
    float aps(altitude) ;
        aps:title = "hybrid pressure at midlayers" ;
        aps:units = "Pa" ;
    float bps(altitude) ;
        bps:title = "hybrid sigma at midlayers" ;
        bps:units = "" ;
    float ap(interlayer) ;
        ap:title = "hybrid pressure at interlayers" ;
        ap:units = "Pa" ;
    float bp(interlayer) ;
        bp:title = "hybrid sigma at interlayers" ;
        bp:units = "" ;
```

```

float soildepth(subsurface_layers) ;
    soildepth:long_name = "Soil mid-layer depth" ;
    soildepth:units = "m" ;
    soildepth:positive = "down" ;
float cu(latitude, rlonu) ;
    cu:title = "Conversion coefficients cov <--> natural" ;
float cv(rlatv, longitude) ;
    cv:title = "Conversion coefficients cov <--> natural" ;
float aire(latitude, longitude) ;
    aire:title = "Mesh area" ;
float phisinit(latitude, longitude) ;
    phisinit:title = "Geopotential at the surface" ;
float emis(Time, latitude, longitude) ;
    emis:title = "Surface emissivity" ;
    emis:units = "w.m-1" ;
float tsurf(Time, latitude, longitude) ;
    tsurf:title = "Surface temperature" ;
    tsurf:units = "K" ;
float ps(Time, latitude, longitude) ;
    ps:title = "surface pressure" ;
    ps:units = "Pa" ;
float co2ice(Time, latitude, longitude) ;
    co2ice:title = "co2 ice thickness" ;
    co2ice:units = "kg.m-2" ;
float mtot(Time, latitude, longitude) ;
    mtot:title = "total mass of water vapor" ;
    mtot:units = "kg/m2" ;
float icetot(Time, latitude, longitude) ;
    icetot:title = "total mass of water ice" ;
    icetot:units = "kg/m2" ;
float tauTES(Time, latitude, longitude) ;
    tauTES:title = "tau abs 825 cm-1" ;
    tauTES:units = "" ;
float h2o_ice_s(Time, latitude, longitude) ;
    h2o_ice_s:title = "surface h2o_ice" ;
    h2o_ice_s:units = "kg.m-2" ;
}

```

The structure of the file is thus as follows:

- the dimensions
- variable “time” containing the time of the timestep stored in the file (in Martian days since the beginning of the run)
- variable “control” containing many parameters, as described above.
- from “ rhonu” to ’phisinit”: a list of data describing the geometrical coordinates of the data file, plus the surface topography
- finally, all the 2D or 3D data stored in the run.

6.3.3 Stats files

As an option (`stats` must be set to `.true.` in `callphys.def`), the model can accumulate any variable from any subroutine of the physics by calling subroutine `wstat`

This save is performed at regular intervals 12 times a day. An average of the daily evolutions over the whole run is calculated (for example, for a 10 day run, the averages of the variable values at 0hTU, 2hTU, 4hTU,...24hTU are calculated), along with RMS standard deviations of the variables. This output is given in file `stats.nc`.

Illustrative example of the contents of a `stats.nc` file (using `ncdump`):
`ncdump -h stats.nc`

```

netcdf stats {
dimensions:
    latitude = 49 ;
    longitude = 65 ;
    altitude = 25 ;
    llmp1 = 26 ;
    Time = UNLIMITED ; // (12 currently)
variables:
    float Time(Time) ;
        Time:title = "Time" ;
        Time:units = "days since 0000-00-0 00:00:00" ;
    float latitude(latitude) ;
        latitude:title = "latitude" ;
        latitude:units = "degrees_north" ;
    float longitude(longitude) ;
        longitude:title = "East longitude" ;
        longitude:units = "degrees_east" ;
    float altitude(altitude) ;
        altitude:long_name = "altitude" ;
        altitude:units = "km" ;
        altitude:positive = "up" ;
    float aps(altitude) ;
        aps:title = "hybrid pressure at midlayers" ;
        aps:units = "" ;
    float bps(altitude) ;
        bps:title = "hybrid sigma at midlayers" ;
        bps:units = "" ;
    float ps(Time, latitude, longitude) ;
        ps:title = "Surface pressure" ;
        ps:units = "Pa" ;
    float ps_sd(Time, latitude, longitude) ;
        ps_sd:title = "Surface pressure total standard deviation over the season" ;
        ps_sd:units = "Pa" ;
    float tsurf(Time, latitude, longitude) ;
        tsurf:title = "Surface temperature" ;
        tsurf:units = "K" ;
    float tsurf_sd(Time, latitude, longitude) ;
        tsurf_sd:title = "Surface temperature total standard deviation over the season" ;
        tsurf_sd:units = "K" ;
    float co2ice(Time, latitude, longitude) ;
        co2ice:title = "CO2 ice cover" ;
        co2ice:units = "kg.m-2" ;
    float co2ice_sd(Time, latitude, longitude) ;
        co2ice_sd:title = "CO2 ice cover total standard deviation over the season" ;
        co2ice_sd:units = "kg.m-2" ;
    float fluxsurf_lw(Time, latitude, longitude) ;
        fluxsurf_lw:title = "Thermal IR radiative flux to surface" ;
        fluxsurf_lw:units = "W.m-2" ;
    float fluxsurf_lw_sd(Time, latitude, longitude) ;
        fluxsurf_lw_sd:title = "Thermal IR radiative flux to surface total standard deviation over the season" ;
        fluxsurf_lw_sd:units = "W.m-2" ;
    float fluxsurf_sw(Time, latitude, longitude) ;
        fluxsurf_sw:title = "Solar radiative flux to surface" ;
        fluxsurf_sw:units = "W.m-2" ;
    float fluxsurf_sw_sd(Time, latitude, longitude) ;
        fluxsurf_sw_sd:title = "Solar radiative flux to surface total standard deviation over the season" ;
        fluxsurf_sw_sd:units = "W.m-2" ;
    float fluxtop_lw(Time, latitude, longitude) ;
        fluxtop_lw:title = "Thermal IR radiative flux to space" ;
        fluxtop_lw:units = "W.m-2" ;
    float fluxtop_lw_sd(Time, latitude, longitude) ;
        fluxtop_lw_sd:title = "Thermal IR radiative flux to space total

```

```

standard deviation over the season" ;
    fluxtop_lw_sd:units = "W.m-2" ;
float fluxtop_sw(Time, latitude, longitude) ;
    fluxtop_sw:title = "Solar radiative flux to space" ;
    fluxtop_sw:units = "W.m-2" ;
float fluxtop_sw_sd(Time, latitude, longitude) ;
    fluxtop_sw_sd:title = "Solar radiative flux to space total standard deviation over the season" ;
    fluxtop_sw_sd:units = "W.m-2" ;
float dod(Time, latitude, longitude) ;
    dod:title = "Dust optical depth" ;
    dod:units = "" ;
float dod_sd(Time, latitude, longitude) ;
    dod_sd:title = "Dust optical depth total standard deviation over the season" ;
    dod_sd:units = "" ;
float temp(Time, altitude, latitude, longitude) ;
    temp:title = "Atmospheric temperature" ;
    temp:units = "K" ;
float temp_sd(Time, altitude, latitude, longitude) ;
    temp_sd:title = "Atmospheric temperature total standard deviation over the season" ;
    temp_sd:units = "K" ;
float u(Time, altitude, latitude, longitude) ;
    u:title = "Zonal (East-West) wind" ;
    u:units = "m.s-1" ;
float u_sd(Time, altitude, latitude, longitude) ;
    u_sd:title = "Zonal (East-West) wind total standard deviation over the season" ;
    u_sd:units = "m.s-1" ;
float v(Time, altitude, latitude, longitude) ;
    v:title = "Meridional (North-South) wind" ;
    v:units = "m.s-1" ;
float v_sd(Time, altitude, latitude, longitude) ;
    v_sd:title = "Meridional (North-South) wind total standard deviation over the season" ;
    v_sd:units = "m.s-1" ;
float w(Time, altitude, latitude, longitude) ;
    w:title = "Vertical (down-up) wind" ;
    w:units = "m.s-1" ;
float w_sd(Time, altitude, latitude, longitude) ;
    w_sd:title = "Vertical (down-up) wind total standard deviation over the season" ;
    w_sd:units = "m.s-1" ;
float rho(Time, altitude, latitude, longitude) ;
    rho:title = "Atmospheric density" ;
    rho:units = "none" ;
float rho_sd(Time, altitude, latitude, longitude) ;
    rho_sd:title = "Atmospheric density total standard deviation over the season" ;
    rho_sd:units = "none" ;
float q2(Time, altitude, latitude, longitude) ;
    q2:title = "Boundary layer eddy kinetic energy" ;
    q2:units = "m2.s-2" ;
float q2_sd(Time, altitude, latitude, longitude) ;
    q2_sd:title = "Boundary layer eddy kinetic energy total standard deviation over the season" ;
    q2_sd:units = "m2.s-2" ;
float vmr_h2ovapor(Time, altitude, latitude, longitude) ;
    vmr_h2ovapor:title = "H2O vapor volume mixing ratio" ;
    vmr_h2ovapor:units = "mol/mol" ;
float vmr_h2ovapor_sd(Time, altitude, latitude, longitude) ;
    vmr_h2ovapor_sd:title = "H2O vapor volume mixing ratio total standard deviation over the season" ;
    vmr_h2ovapor_sd:units = "mol/mol" ;
float vmr_h2oice(Time, altitude, latitude, longitude) ;
    vmr_h2oice:title = "H2O ice volume mixing ratio" ;

```

```

        vmr_h2o_ice:units = "mol/mol" ;
        float vmr_h2o_ice_sd(Time, altitude, latitude, longitude) ;
        vmr_h2o_ice_sd:title = "H2O ice volume mixing ratio total standard deviation over the season" ;
        vmr_h2o_ice_sd:units = "mol/mol" ;
        float mtot(Time, latitude, longitude) ;
        mtot:title = "total mass of water vapor" ;
        mtot:units = "kg/m2" ;
        float mtot_sd(Time, latitude, longitude) ;
        mtot_sd:title = "total mass of water vapor total standard deviation over the season" ;
        mtot_sd:units = "kg/m2" ;
        float icetot(Time, latitude, longitude) ;
        icetot:title = "total mass of water ice" ;
        icetot:units = "kg/m2" ;
        float icetot_sd(Time, latitude, longitude) ;
        icetot_sd:title = "total mass of water ice total standard deviation over the season" ;
        icetot_sd:units = "kg/m2" ;
    }

```

The structure of the file is similar to the `diagfi.nc` file, except that, as stated before, the average of variables are given for 12 times of the day and that RMS standard deviation are also provided.

Chapter 7

Zoomed simulations

The LMD GCM can use a zoom to enhance the resolution locally. In practice, one can increase the latitudinal resolution on the one hand, and the longitudinal resolution on the other hand.

7.1 To define the zoomed area

The zoom is defined in `run.def`. Here are the variables that you want to set:

- East longitude (in degrees) of zoom center `clon`
- latitude (in degrees) of zoom center `clat`
- zooming factors, along longitude `grossismx`. *Typically 1.5, 2 or even 3 (see below)*
- zooming factors, along latitude `grossismy`. *Typically 1.5, 2 or even 3 (see below)*
- `fxyhyphb`: **must be set to "T" for a zoom**, whereas it must be *F* otherwise
- extension in longitude of zoomed area `dzoomx`. This is the total longitudinal extension of the zoomed region (degree).
It is recommended that $grossismx \times dzoomx < 200^\circ$
- extension in latitude of the zoomed region `dzoomy`. This is the total latitudinal extension of the zoomed region (degree).
It is recommended that $grossismy \times dzoomy < 100^\circ$
- stiffness of the zoom along longitudes `taux`. 2 is for a smooth transition in longitude, more means sharper transition.
- stiffness of the zoom along latitudes `tauy`. 2 is for a smooth transition in latitude, more means sharper transition.

7.2 Making a zoomed initial state

One must start from an initial state archive `start_archive.nc` obtained from a previous simulation (see section 4.10) Then compile and run `newstart.e` **using the `run.def` file designed for the zoom.**

After running `newstart.e`. The zoomed grid may be visualized using `grads`, for instance. Here is a `grads` script that can be used to map the grid above a topography map:


```

set mpdraw off
set grid off
sdfopen restart.nc
set gxout grid
set digsiz 0
set lon -180 180
d ps
close 1
*** replace the path to surface.nc in the following line:
sdfopen /u/forget/WWW/datagcm/datafile/surface.nc
set lon -180 180
set gxout contour
set clab off
set cint 3
d zMOL

```

7.3 Running a zoomed simulation and stability issue

- **dynamical timestep** Because of their higher resolution, zoomed simulation requires a higher timestep. Therefore in `run.def`, the number of dynamical timestep per day `day_step` must be increased by more than `grossismx` or `grossismy` (twice that if necessary). However, you can keep the same physical timestep (48/sol) and thus increase `iphysiq` accordingly (`iphysiq = day_step/48`).
- It has been found that when zooming in longitude, one must set `ngroup=1` in `dyn3d/groupeun.F`. Otherwise the run is less stable.
- The very first initial state made with `newstart.e` can be noisy and dynamically unstable. It may be necessary to strongly increase the intensity of the dissipation and increase `day_step` in `run.def` for 1 to 3 sols, and then use less strict values.
- If the run remains very unstable and requires too much dissipation or a too small timestep, a good tip to help stabilize the model is to decrease the vertical extension of your run and the number of layer (one generally zoom to study near-surface process, so 20 to 22 layers and a vertical extension up to 60 or 80 km is usually enough).

Chapter 8

Water Cycle Simulation

In order to simulate the water cycle with the LMD GCM:

- In `callphys.def`, set tracer to true: `tracer=.true..`
- It is best to run with a semi-interactive dust (in order to better represent the evolution available condensation nuclei) and thus set `dustbin = 2`, and use corresponding additional tracers.
- Use the same options as shown below for the Tracer part, the rest does not change compared to the basic `callphys.def`.

```
## Tracer (dust water, ice and/or chemical species) options (used if tracer=T):
## ~~~~~
# DUST: Transported dust ? (if >0, use 'dustbin' dust bins)
dustbin = 2
# DUST: Radiatively active dust ? (matters if dustbin>0)
active = .true.
# DUST: use mass and number mixing ratios to predict dust size ?
doubleq = .true.
# DUST: lifted by GCM surface winds ?
lifting = .true.
# DUST: lifted by dust devils ?
calldevil = .false.
# DUST: Scavenging by H2O/CO2 snowfall ?
scavenging = .true.
# DUST/WATERICE: Gravitationnal sedimentation ?
sedimentation = .true.
# WATERICE: Radiatively active transported atmospheric water ice ?
activice = .true.
# WATER: Compute water cycle
water = .true.
# WATER: Microphysical scheme for water-ice clouds?
microphys = .true.
# WATER: parameter contact
mteta = 0.95
# WATER: current permanent caps at both poles. True IS RECOMMENDED
#         (with .true., North cap is a source of water and South pole
#         is a cold trap)
caps = .true.
# WATER: Water ice albedo ?
albedo_h2o_ice = 0.35
# WATER: Water ice thermal inertia
inert_h2o_ice = 800
# WATER: Frost thickness threshold for albedo
frost_albedo_threshold = 0.005
nuice_sed=0.1
icedryness=1
```

- **Settings** You need to run with at least 4 tracers (if you don't have dust (`dustbin=0`) or other chemical species (`photochem=F`). An appropriate `tracer.def` for running with `co2`, dust and water cycle should be

```
7
co2
dust_mass
dust_number
h2o_vap
h2o_ice
ccn_mass
ccn_number
```

Note that the order in which tracers are set in the `tracer.def` file is not important.

- **Run**

Same as usual. Just make sure that your start files contains the initial states for all the tracers you use (or else compile and run **newstart.e** to initialize them).

Chapter 9

Photochemical Module

The LMD GCM now includes a photochemical module, which allows to compute the atmospheric composition.

- 14 chemical species are included: CO₂ (background gas), CO, O, O(¹D), O₂, O₃, H, H₂, OH, HO₂, H₂O₂, N₂, Ar (inert) and H₂O.
- In `callphys.def`, set tracer to true `tracer=.true..` Use the same options as shown below for the tracer part of `callphys.def`. You need to set `photochem=.true.`, and to include the water cycle (`water=.true.`, `sedimentation=.true.`; see Chapter 8), because composition is extremely dependent on the water vapor abundance.
- You will need the up-to-date file `jmars.yyyymmdd` (e.g. `jmars.20030707`), which contains the photodissociation rates. It should be in the `datafile` directory in which are stored datafiles used by the GCM (the path to these files is set in file `datafile.h`, in the `phymars` directory).

- **Settings**

Compile as usual, what really matters are the options in `callphys.def` and `traceur.def`. You need at least 19 tracers: 13 chemical species (`co2`, `co`, `o`, `o(1d)`, `o2`, `o3`, `h`, `h2`, `oh`, `ho2`, `h2o2`, `n2`, `ar`) along with water vapor (`h2o_vap`), water ice (`h2o_ice`), related condensation nuclei moments (`ccn_mass` and `ccn_number`), and dust moments (`dust_mass` and `dust_number`).

the `traceur.def` file should thus be something like:

```
19
co2
co
o
o1d
o2
o3
h
h2
oh
ho2
h2o2
n2
ar
```

dust_mass
dust_number
h2o_vap
h2o_ice
ccn_mass
ccn_number

- **Run**

Same as usual. Just make sure that your start files contains the correct number of tracers. If you need to initialize the composition, you can run **newstart** and use the options

- ini_q: the 15 tracers are initialized, including water ice and vapor.
- ini_q-h2o: the 13 chemical species are initialized, water ice is put to zero, and water vapor is kept untouched.
- ini_q-iceh2o: the 13 chemical species are initialized, water ice and vapor are kept untouched.

The initialization is done with the files `atmosfera_LMD_may.dat` and `atmosfera_LMD_min.dat`, which should also be found in the *datafile* directory.

- **Outputs**

The outputs can be done from the `aeronomars/calchim.F` routine for the 14 chemical species. The variables put in the `diagfi.nc` and `stats.nc` files are labeled (where *name* is the name of the chemical species, e.g. `co2`):

- *n_name*: local density (in molecule cm^{-3} , 3-dimensional field)
- *c_name*: integrated column density (in molecule cm^{-2} , 2-dimensional field)

Chapter 10

1D version of the Mars model

The physical part of the model can be used to generate realistic 1-D simulations (one atmosphere column). In practice, the simulation is controlled from a main program called `testphys1d.F` which, after initialization, then calls the master subroutine of the physics `physiq.F` described in the preceding chapters.

10.1 Compilation

To compile the Martian model in 1-D with 29 layers, simply type (in compliance with the `makelmdz_fcm` function manual described in section 5.4)

```
makelmdz_fcm -arch local -d 29 -p mars testphys1d
```

You will find executable `testphys1d_29_phymars_seq.e` (the compiled model) in the `bin` subdirectory from which you ran the `makelmdz_fcm` command.

10.2 1-D runs and input files

The 1-D model does not use an initial state file (the simulation must be long enough to obtain a balanced state). Thus, to generate a simulation simply type:

```
./testphys1d.e
```

The following example files are available in the `deftank` directory (copy them into your working directory first):

- `callphys.def` : controls the options in the physics, just like for the 3D GCM.
- `z2sig.def` : controls the vertical discretization (no change needed, in general), functions as with the 3D GCM.
- `traceur.def` : controls the tracer names (this file may not be present, as long as you run without tracers (option `tracer=.false.` in `callphys.def`))
- `run.def` : controls the 1-D run parameters and initializations (this is actually file `run.def.1d` the `deftank` directory, which must be renamed `run.def` to be read by the program).

The last file is different from the 3D GCM's `run.def` input file, as it contains options specific to the 1-D model, as shown in the example below:

```
#
#-----
# Run parameters for the 1D 'testphys1d.e' model
#-----
```

```

##### Time integration parameters
#
# Initial date (in martian sols ; =0 at Ls=0)
day0=0
# Initial local time (in hours, between 0 and 24)
time=0
# Number of time steps per sol
day_step=48
# Number of sols to run
ndt = 100

##### Physical parameters
#
# Surface pressure (Pa)
psurf= 610
# Reference dust opacity at 700 Pa, in the visible (true tau~tauref*psurf/700)
tauvis=0.2
# latitude (in degrees)
latitude= 0.
# Albedo of bare ground
albedo=0.2
# Soil thermal inertia (SI)
inertia=400
# zonal eastward component of the geostrophic wind (m/s)
u=10.
# meridional northward component of the geostrophic wind (m/s)
v=0.
# Initial CO2 ice on the surface (kg.m-2)
co2ice=0
# hybrid vertical coordinate ? (.true. for hybrid and .false. for sigma levels)
hybrid=.true.

##### Initial atmospheric temperature profile
#
# Type of initial temperature profile
#   ichoice=1   Constant Temperature: T=tref
#   ichoice=2   Savidjari profile (as Seiff but with dT/dz=cte)
#   ichoice=3   Lindner (polar profile)
#   ichoice=4   inversion
#   ichoice=5   Seiff (standard profile, based on Viking entry)
#   ichoice=6   constant T + gaussian perturbation (levels)
#   ichoice=7   constant T + gaussian perturbation (km)
#   ichoice=8   Read in an ascii file "profile"
ichoice=5
# Reference temperature tref (K)
tref=200
# Add a perturbation to profile if isin=1
isin=0
# peak of gaussian perturbation (for ichoice=6 or 7)
pic=26.522
# width of the gaussian perturbation (for ichoice=6 or 7)
largeur=10
# height of the gaussian perturbation (for ichoice=6 or 7)
hauteur=30.

# some definitions for the physics, in file 'callphys.def'
INCLUDEDEF=callphys.def

```

Note that, just as for the 3-D GCM run.def file, input parameters may be given in any order, or even not given at all (in which case default values are used by the program).

Initial conditions for tracers can be provided as ASCII text files; when **testphys1d.e** runs, it looks for a `profile_*` file, where `*` stands for a tracer name (as given in the `traceur.def` file) to initialize the tracer profile. The data in the file should be given as one

value per line, first line for the surface value (in kg/m²), followed by values for each atmospheric layer (going from bottom to top of the atmosphere). If no `profile.*` file is found, then the tracer values are initialized to zero (except for the 'co2' tracer, which will be set to 0.95).

10.3 Output data

During the entire 1D simulation, you can obtain output data for any variable from any physical subroutine by using subroutine `writediagfi` (as when running 3D simulations) or `writegld`. This subroutine creates file `gld.nc` that can be read by GRADS. This subroutine is typically called at the end of subroutine `physiq`.

Example of a call to subroutine `writegld` requesting temperature output: (`ngrid` horizontal point, `nlayer` layers, variable `pt` called "T" in K units):

```
CALL writegld(ngrid,nlayer,pt,'T','K')
```


Appendix A

GCM Martian Calendar

For Mars, dates and seasons are expressed in Solar Longitude (L_s , in degrees or in radians) counting from the northern hemisphere spring equinox. In the GCM, time is counted in Martian solar days, or “sols” (1 sols = 88775 s) from the northern spring equinox. The following table gives the correspondence between sols and L_s , calculated for the GCM using one Martian year = 669 sols exactly.

sol	L_s		sol	L_s		sol	L_s		
0.	360.000	Spring equinox N	240.	111.455		480.	247.408		
5.	2.550		245.	113.816		485.	250.666		
10.	5.080		250.	116.190		490.	253.925		
15.	7.590		255.	118.578		495.	257.182		
20.	10.081		260.	120.981		500.	260.435		
25.	12.554		265.	123.400		505.	263.683		
30.	15.009		270.	125.835		510.	266.924		
						514.76	270.		Winter solstice N
35.	17.447		275.	128.287		515.	270.156		
40.	19.869		280.	130.756		520.	273.377		
45.	22.275		285.	133.243		525.	276.587		
50.	24.666		290.	135.750		530.	279.783		
55.	27.043		295.	138.275		535.	282.965		
60.	29.407	300.	140.821	540.	286.130				
65.	31.758	305.	143.388	545.	289.277				
70.	34.096	310.	145.975	550.	292.406				
75.	36.423	315.	148.585	555.	295.515				
80.	38.739	320.	151.217	560.	298.604				
85.	41.046	325.	153.872	565.	301.671				
90.	43.343	330.	156.550	570.	304.715				
95.	45.631	335.	159.251	575.	307.737				
100.	47.912	340.	161.977	580.	310.735				
105.	50.186	345.	164.727	585.	313.709				
110.	52.453	350.	167.502	590.	316.658				
115.	54.714	355.	170.301	595.	319.583				
120.	56.970	360.	173.126	600.	322.483				
125.	59.222	365.	175.975	605.	325.358				
130.	61.471	370.	178.850	610.	328.207				
		371.99	180.	Autumn equinox N					
135.	63.716	375.	181.750	615.	331.032				
140.	65.959	380.	184.675	620.	333.831				
145.	68.201	385.	187.624	625.	336.606				
150.	70.442	390.	190.598	630.	339.356				
155.	72.683	395.	193.596	635.	342.082				
160.	74.925	400.	196.618	640.	344.783				
165.	77.168	405.	199.662	645.	347.461				
170.	79.413	410.	202.729	650.	350.116				
175.	81.661	415.	205.818	655.	352.748				
180.	83.912	420.	208.927	660.	355.357				
185.	86.167	425.	212.056	665.	357.945				
				669.	0.	Spring equinox N			
190.	88.427	430.	215.203	670.	0.512				
193.47	90.	Summer solstice N							
195.	90.693	435.	218.368	675.	3.057				
200.	92.965	440.	221.549	680.	5.583				
205.	95.245	445.	224.746	685.	8.089				
210.	97.532	450.	227.955	690.	10.577				
215.	99.827	455.	231.177	695.	13.046				
220.	102.131	460.	234.409	700.	15.498				
225.	104.446	465.	237.650	705.	17.933				
230.	106.770	470.	240.898	710.	20.351				
235.	109.107	475.	244.151	715.	22.755				
240.	111.455	480.	247.408	720.	25.143				

Appendix B

Utilities

A few post-processing tools, which handle GCM outputs (files `diagfi.nc` and `stats.nc`) are available in the `LMDZ.MARS/util` subdirectory. This directory contains the sources codes along with a `README.exec` file which explains what the various programs are for and how to compile them.

B.1 `concatnc`

This program concatenates consecutive output files (`diagfi.nc` or even `stats.nc` files) for a selection of variable, in order to obtain one single big file. The time dimension of the output can be "sols" or "Ls" (note that in that latter case, Ls values won't be evenly distributed, and software like Grads may not be able to use and plot the data).

To obtain an evenly sampled "Ls" timescale, you can use the `lslin.e` program (described below). The output file created by `concatnc.e` is `concat.nc`

B.2 `lslin`

This program is designed to interpolate data given in irregular Solar Longitude (Ls) into an evenly sampled linear time coordinate (usable with Grads). Input Netcdf files may be `diagfi.nc` or `concat.nc` files and the resulting output file is `lslin.nc` `lslin` also create a `lslin.ct1` file that can be read directly by grads (`>xdfopen lslin.ct1`) to plot in Ls coordinate to avoid some problem with grads when Grads think that "the time interval is too small"...

B.3 `localtime`

The `localtime.e` program is designed to re-interpolate data in order to yield values at the same given local time (useful to mimic satellite observations, or analyse day to day variations at given local time).

Input files may be of `diagfi.nc`, `stats.nc` or `concat.nc` type and the output file name is build from the input one, to which `_LT.nc` is appened (e.g. if the input file is `myfile.nc` then output file will be `myfile_LT.nc`).

B.4 `zrecast`

With this program you can recast atmospheric (i.e.: 4D-dimensional longitude-latitude-altitude-time) data from GCM outputs (e.g. as given in `diagfi.nc`, `concat.nc` and `stats.nc` files) onto either *pressure* or *altitude above areoid* vertical coordinates.

Since integrating the hydrostatic equation is required to recast the data, the input file must contain surface pressure and atmospheric temperature, as well as the ground geopotential.

If recasting data onto *pressure* coordinates, then the output file name is given by the input file name to which `_P.nc` will be appened. If recasting data onto *altitude above areoid* coordinates, then a `_A.nc` will be appened.

B.5 `lslin`

This program is designed to interpolate data in Solar Longitude (Ls) linear time coordinate (usable with grads) `diagfi.nc` or `concat.nc` files.

lslin also create a lslin.ct1 file that can be read directly by grads (>xdfopen lslin.ct1) to plot in Ls coordinate to avoid some problem with grads when grads think that "the time interval is too small"...

B.6 hrecast

This program can interpolate GCM output on any horizontal grid (regular lat - lon) as long as it cover all the planet. Useful to compare runs obtained at different horizontal resolutions.

B.7 expandstartfi

This program takes a physics start file (startfi.nc) and recasts it on the corresponding lonxlat grid (so its contents may easily be displayed using Grads, Ferret, etc.)

B.8 extract

This program extracts (ie: interpolates) pointwise values of an atmospheric variable from a 'zrecast'ed diagfi file (works if altitude is geometrical height or a pressure vertical coordinates).

Bibliography

- [1] M. Angelats i Coll, F. Forget, M. A. López-Valverde, and F. González-Galindo. The first Mars thermospheric general circulation model: The Martian atmosphere from the ground to 240 km. *Geophys. Res. Lett.*, 32:4201, 2005.
- [2] J.-L. Dufresne, R. Fournier, C. Hourdin, and F. Hourdin. Net Exchange Reformulation of Radiative Transfer in the CO₂ 15- μ m Band on Mars. *Journal of Atmospheric Sciences*, 62:3303–3319, 2005.
- [3] F. Forget, F. Hourdin, R. Fournier, C. Hourdin, O. Talagrand, M. Collins, S. R. Lewis, P. L. Read, and J.-P. Huot. Improved general circulation models of the Martian atmosphere from the surface to above 80 km. *J. Geophys. Res.*, 104:24,155–24,176, 1999.
- [4] F. Forget, F. Hourdin, and O. Talagrand. CO₂ snow fall on Mars: Simulation with a general circulation model. *Icarus*, 131:302–316, 1998.
- [5] F. González-Galindo, F. Forget, M. A. López-Valverde, and M. Angelats i Coll. A Ground-to-Exosphere Martian General Circulation Model. 2. The Atmosphere During Perihelion Conditions: Thermospheric Polar Warming. *Journal of Geophysical Research (Planets)*, 114(E13):E08004, 2009.
- [6] F. González-Galindo, F. Forget, M. A. López-Valverde, M. Angelats i Coll, and E. Millour. A Ground-to-Exosphere Martian General Circulation Model. 1. Seasonal, Diurnal and Solar Cycle Variation of Thermospheric Temperatures. *Journal of Geophysical Research (Planets)*, 114(E13):4001, 2009.
- [7] F. González-Galindo, M. A. López-Valverde, M. Angelats i Coll, and F. Forget. Extension of a Martian general circulation model to thermospheric altitudes: UV heating and photochemical models. *Journal of Geophysical Research (Planets)*, 110(E9):9008, 2005.
- [8] C. Hourdin, J.-L. Dufresnes, R. Fournier, and F. Hourdin. Net exchange reformulation of radiative transfer in the CO₂ 15 μ m band on mars. Article in preparation, 2000.
- [9] F. Hourdin. A new representation of the CO₂ 15 μ m band for a Martian general circulation model. *J. Geophys. Res.*, 97(E11):18,319–18,335, 1992.
- [10] F. Hourdin and A. Armengaud. Test of a hierarchy of finite-volume schemes for transport of trace species in an atmospheric general circulation model. *Mon. Wea. Rev.*, 127:822–837, 1999.
- [11] F. Hourdin, P. Le Van, F. Forget, and O. Talagrand. Meteorological variability and the annual surface pressure cycle on Mars. *J. Atmos. Sci.*, 50:3625–3640, 1993.
- [12] F. Lefèvre, J.-L. Bertaux, R. T. Clancy, T. Encrenaz, K. Fast, F. Forget, S. Lebonnois, F. Montmessin, and S. Perrier. Heterogeneous chemistry in the atmosphere of Mars. *Nature*, 454:971–975, 2008.
- [13] F. Lefèvre, S. Lebonnois, F. Montmessin, and F. Forget. Three-dimensional modeling of ozone on Mars. *Journal of Geophysical Research (Planets)*, 109:E07004, 2004.
- [14] S. R. Lewis, M. Collins, P. L. Read, F. Forget, F. Hourdin, R. Fournier, C. Hourdin, O. Talagrand, and J.-P. Huot. A climate database for Mars. *J. Geophys. Res.*, 104:24,177–24,194, 1999.
- [15] F. Lott and M. Miller. A new sub-grid scale orographic drag parametrization: its formulation and testing. *Q. J. R. Meteorol. Soc.*, 123:101–128, 1997.
- [16] J.-B. Madeleine, F. Forget, E. Millour, L. Montabone, and M. J. Wolff. Revisiting the radiative impact of dust on Mars using the LMD Global Climate Model. *Journal of Geophysical Research (Planets)*, 116:11010, Nov. 2011.
- [17] F. Montmessin, F. Forget, P. Rannou, M. Cabane, and R. M. Haberle. Origin and role of water ice clouds in the Martian water cycle as inferred from a general circulation model. *Journal of Geophysical Research (Planets)*, 109(E18):10004, 2004.

- [18] O. B. Toon, C. P. McKay, T. P. Ackerman, and K. Santhanam. Rapid calculation of radiative heating rates and photodissociation rates in inhomogeneous multiple scattering atmospheres. *J. Geophys. Res.*, 94:16,287–16,301, 1989.
- [19] M. J. Wolff, M. D. Smith, R. T. Clancy, R. Arvidson, M. Kahre, F. Seelos, S. Murchie, and H. Savijärvi. Wavelength dependence of dust aerosol single scattering albedo as observed by the Compact Reconnaissance Imaging Spectrometer. *Journal of Geophysical Research (Planets)*, 114(E13):0–+, 2009.